

- Q1 Q0 A simple pendulum of mass $m=20$ kg and length L is pulled
Q0 back and held with a horizontal force of 100 N (see Fig 1).
Q0 The tension in the string at this equilibrium position is:
Q0
A1 220 N
A2 60 N
A3 120 N
A4 190 N
A5 260 N
Q0
- Q2 Q0 A horizontal aluminum rod (shear modulus = 2.5×10^{10} N/m²)
Q0 projects $L=5.0$ cm from the wall (see Fig 6). The cross sectional
Q0 area of the rod $A = 1.0 \times 10^{-5}$ m². A shearing force of 500 N
Q0 is applied at the end of the rod. Find the vertical deflection
Q0 $\Delta(x)$ of the end of the rod.
Q0
A1 1.0×10^{-4} m
A2 2.0×10^{-4} m
A3 3.0×10^{-4} m
A4 4.0×10^{-4} m
A5 5.0×10^{-4} m
Q0
- Q3 Q0 A uniform rod AB is 1.2 m long and weighs 16 N. It is suspended
Q0 by strings AC and BD as shown in Fig 2. A block P weighing 96 N
Q0 is attached at point E, 0.30 m from A. The tension in the string
Q0 BD is:
Q0
A1 32 N
A2 24 N
A3 64 N
A4 48 N
A5 112 N
Q0
- Q4 Q0 Four point masses are at the corners of a square whose side is
Q0 20 cm long (see Fig 3). What is the magnitude of the net
Q0 gravitational force on a point mass $m_5 = 2.5$ kg located at the
Q0 center of the square?
Q0
A1 3.3×10^{-8} N
A2 1.1×10^{-8} N
A3 2.2×10^{-8} N
A4 4.4×10^{-8} N
A5 6.6×10^{-8} N
Q0
- Q5 Q0 An object is fired vertically upward from the surface of
Q0 the Earth (Radius = R) with an initial speed of $(V_{esc})/2$,
Q0 where (V_{esc} = escape speed). Neglecting air resistance,
Q0 how far above the surface of Earth will it reach?
Q0
A1 $R/3$
A2 $R/2$
A3 $3R$
A4 $2R$
A5 R
Q0




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9

$$M_{\text{earth}} = 5.98 \times 10^{24} \text{ kg}$$

$$R_{\text{earth}} = 6.37 \times 10^6 \text{ m}$$

$$G = 6.67 \times 10^{-11} \text{ N}\cdot\text{m}^2/\text{kg}^2$$

- Q6 Q0 What is the escape speed on a spherical planet whose radius
Q0 is 3200 km and whose gravitational acceleration at the surface
Q0 is 4.00 m/s^2 ?
Q0
A1 5.06 km/s
A2 3.58 km/s
A3 11.2 km/s
A4 9.80 km/s
A5 4.00 km/s
Q0
- Q7 Q0 A planet requires 300 (Earth) days to complete its circular
Q0 orbit about its sun (mass $M = 6.0 \times 10^{30} \text{ kg}$).
Q0 The orbital speed of the planet is:
Q0
A1 $4.6 \times 10^4 \text{ m/s}$
A2 $5.4 \times 10^4 \text{ m/s}$
A3 $6.5 \times 10^4 \text{ m/s}$
A4 $3.5 \times 10^4 \text{ m/s}$
A5 $7.5 \times 10^4 \text{ m/s}$
Q0
- Q8 Q0 A water hose of 1.00 cm radius is used to fill a container of
Q0 volume $20.0 \times 10^3 \text{ cm}^3$. It takes 60 s to fill the container.
Q0 What is the speed at which the water leaves the hose?
Q0
A1 106 cm/s
A2 201 cm/s
A3 154 cm/s
A4 189 cm/s
A5 255 cm/s
Q0
- Q9 Q0 Water enters a house through a pipe with a velocity of 4.0 m/s
Q0 at a pressure of $4 \times 10^5 \text{ Pa}$. The water in a narrower pipe
Q0 at the second floor bathroom 5.0 m above has a velocity of
Q0 16 m/s. What is the pressure of water in the bathroom?
Q0 (Density of water = $1.0 \times 10^3 \text{ kg/m}^3$)
Q0
A1 $2.3 \times 10^5 \text{ Pa}$
A2 $1.5 \times 10^5 \text{ Pa}$
A3 $5.5 \times 10^5 \text{ Pa}$
A4 $4.5 \times 10^5 \text{ Pa}$
A5 $3.0 \times 10^5 \text{ Pa}$
Q0
- Q10 Q0 A block of metal has mass of 0.50 kg and density of 8.0×10^3
Q0 kg/m^3 . It is suspended from a string and completely
Q0 submerged in water. Find the tension in the string.
Q0 (Density of water = $1.0 \times 10^3 \text{ kg/m}^3$)
Q0
A1 4.3 N
A2 5.0 N
A3 0.60 N
A4 4.9 N
A5 5.5 N
Q0
- Q11 Q0 A piston of radius $R_1 = 5.0 \text{ cm}$ is used in a hydraulic press to
Q0 exert a force F_1 on the enclosed liquid to raise a car of weight
Q0 $F_2 = 13,500 \text{ N}$ (see Fig 4). If the radius of the larger piston is
Q0 $R_2 = 15 \text{ cm}$, Find F_1 .
Q0
A1 $1.5 \times 10^3 \text{ N}$
A2 $2.5 \times 10^3 \text{ N}$

- A3 3.5×10^3 N
- A4 4.0×10^3 N
- A5 2.0×10^3 N

Q0

Q12Q0 A block of mass 0.50 kg is attached to a horizontal spring (k = 160 N/m). The block is pulled a distance 20 cm from its unstretched position on a frictionless horizontal surface. What is the magnitude of its maximum acceleration?

Q0

- A1 64 m/s²
- A2 0.80 m/s²
- A3 0.28 m/s²
- A4 72 m/s²
- A5 1.9 m/s²

Q0

Q13Q0 A simple pendulum of length = L1 on Earth oscillates with a period = T. Another pendulum of length = L2 on the Moon oscillates with a period = 2*T. Find the ratio L1/L2. (Take g on Moon = (1/6)*g on Earth.)

Q0

- A1 3/2
- A2 1/2
- A3 1/4
- A4 2/3
- A5 2

Q0

Q14Q0 A block-spring system has an amplitude of 4.0 cm and a maximum speed of 0.60 m/s. What is the frequency of oscillation?

Q0

- A1 2.39 Hz
- A2 120 Hz
- A3 60 Hz
- A4 240 Hz
- A5 0.50 Hz

Q0

Q15Q0 A particle oscillates according to the equation:

Q0 $x = 0.20 \cdot \cos(\pi t)$, where $\pi = 3.14$.

Q0 What is the period of the motion?

Q0

- A1 2.0 s
- A2 2.0 Hz
- A3 0.20 s
- A4 π s
- A5 1.0 s

Q0

Q16Q0 A ball (mass=m) is dropped from a bridge that is 40 m high (see Fig 9). It falls directly into a boat, moving with constant velocity, that is 12 m from the point of impact when the ball is released. What is the speed (v) of the boat?

Q0

- A1 4.2 m/s
- A2 10 m/s
- A3 7.4 m/s
- A4 2.5 m/s
- A5 9.5 m/s

Q0

Q17Q0 If $A = 3i - 2j$ and $B = 2j$ what is $(A \times B) \cdot B$?

Q0

- A1 0
- A2 12

- A3 4
- A4 -4
- A5 $6i - 4j$

Q18Q0 A player kicks a ball with a velocity of 50.0 m/s at an angle of 30 degrees above the horizontal. Find the time the ball takes to reach the maximum height.

- A1 2.55 s
- A2 1.35 s
- A3 2.00 s
- A4 1.00 s
- A5 5.10 s

Q19Q0 A man of mass 70.0 kg stands on a scale in an elevator. What does the scale read when the elevator accelerates downward at 1.20 m/s^2 ?

- A1 602 N
- A2 770 N
- A3 686 N
- A4 84 N
- A5 980 N

Q20Q0 A box slides down a 30 degree incline. If the coefficient of kinetic friction between the box and the surface of the incline is 0.30. What is the acceleration of the box?

- A1 2.35 m/s^2
- A2 6.96 m/s^2
- A3 4.90 m/s^2
- A4 0 m/s^2
- A5 9.80 m/s^2

Q21Q0 A 4.0 kg cart starts up an incline with a speed of 3.0 m/s and comes to rest 2.0 m up the incline. The total work done on the cart is:

- A1 -18 J
- A2 8.0 J
- A3 12 J
- A4 -4.0 J
- A5 0 J

Q22Q0 A force of 100 N holds an ideal spring having 200 N/m spring constant in compression. The potential energy stored in the spring is:

- A1 25 J
- A2 0.5 J
- A3 5.0 J
- A4 10 J
- A5 200 J

Q23Q0 A 6.0 kg block is released from rest 80 m above the ground. When it has fallen 60 m its kinetic energy is:

- A1 3500 J
- A2 4800 J
- A3 1200 J

A4 120 J

A5 60 J

Q0

Q24Q0 A ball is thrown into the air. As it rises, there is an increase
Q0 in its:

Q0

A1 potential energy

A2 speed

A3 kinetic energy

A4 acceleration

A5 momentum

Q0

Q25Q0 A 10 g bullet is fired horizontally into a 2.0 kg pendulum block
Q0 at rest. The bullet remains embedded in the block and the block
Q0 with the bullet inside rises to a height of 10 cm. What is the
Q0 initial speed (v) of the bullet? (See Fig 5)

Q0

A1 281 m/s

A2 302 m/s

A3 182 m/s

A4 102 m/s

A5 252 m/s

Q0

Q26Q0 A 2.0 kg and 3.0 kg masses are moving along the x-axis. At a
Q0 particular instant, the 2.0 kg has a velocity of 3.0 m/s and
Q0 the 3.0 kg has a velocity of -1.0 m/s. What is the velocity of
Q0 their center of mass?

Q0

A1 0.60 m/s

A2 1.8 m/s

A3 -0.60 m/s

A4 -1.8 m/s

A5 0.00 m/s

Q0

Q27Q0 A cylinder is 0.10 m in radius and its rotational inertia, about
Q0 the axis through O, is $0.020 \text{ kg}\cdot\text{m}^2$. A string is wound around
Q0 the cylinder and pulled with a force of 1.0 N. The angular
Q0 acceleration of the cylinder is (see Fig 7):

Q0

A1 5.0 rad/s^2

A2 10 rad/s^2

A3 15 rad/s^2

A4 20 rad/s^2

A5 2.5 rad/s^2

Q0

Q28Q0 A wheel initially has an angular velocity of 18 rad/s but it is
Q0 slowing at a rate of 2.0 rad/s^2 . By the time it stops it will
Q0 have turned through:

Q0

A1 13 rev

A2 26 rev

A3 39 rev

A4 52 rev

A5 65 rev

Q0

Q29Q0 Three particles, of mass of m , $2m$ and $3m$, are fastened
Q0 to each other and to a rotation axis at O by three massless
Q0 rods, of lengths a , $2a$ and $3a$ respectively (see Fig 8).
Q0 The combination rotates around the rotational axis with
Q0 angular velocity of w . What is the total angular momentum

Q0 of the three particles relative to point O?

Q0

A1 127 $m \cdot w \cdot a^{**2}$

A2 97 $m \cdot w \cdot a^{**2}$

A3 117 $m \cdot w \cdot a^{**2}$

A4 137 $m \cdot w \cdot a^{**2}$

A5 147 $m \cdot w \cdot a^{**2}$

Q0

Q30Q0 When a man on a frictionless rotating seat extends his arms

Q0 horizontally, his rotational kinetic energy:

Q0

A1 must decrease

A2 must increase

A3 must remain the same

A4 may increase or decrease depending on his initial

A4 angular velocity

A5 may increase or decrease depending on his gravitational

A5 potential energy