

Q1.

A ball is thrown straight up and is caught 2.00 s later at the same point. The initial speed of the ball is:

- A) 9.80 m/s
- B) 7.40 m/s
- C) 4.90 m/s
- D) 12.6 m/s
- E) 19.6 m/s

---

Q2.

Two points A and B in the x-y plane, A has the coordinates (0 m, 3 m) and B has the coordinates (4 m, 0 m). The displacement vector that goes from A to B is:

- A)  $(4\hat{i} - 3\hat{j})$  m
- B)  $(3\hat{i} - 4\hat{j})$  m
- C)  $(4\hat{i} + 3\hat{j})$  m
- D)  $(-4\hat{i} - 3\hat{j})$  m
- E)  $(3\hat{i} + 4\hat{j})$  m

---

Q3.

A projectile is fired from the ground with an initial velocity of  $\vec{v}_o = (3.0\hat{i} + 4.0\hat{j})$  m/s . Find the velocity of the projectile just before hitting the ground.

- A)  $\vec{v} = (3.0\hat{i} - 4.0\hat{j})$  m/s
- B)  $\vec{v} = (-3.0\hat{i} + 4.0\hat{j})$  m/s
- C)  $\vec{v} = (-3.0\hat{i} - 4.0\hat{j})$  m/s
- D)  $\vec{v} = (3.0\hat{i} + 4.0\hat{j})$  m/s
- E)  $\vec{v} = (5.0)$  m/s

---

Q4.

Snow is falling vertically at a constant speed of 8.00 m/s relative to the ground. To a driver of a car (travelling horizontally), the snow appears to be falling at an angle of  $60.0^\circ$  from the vertical direction. What is the speed of the car relative to the ground?


- A) 13.9 m/s
- B) 8.00 m/s
- C) 4.00 m/s
- D) 6.93 m/s
- E) 10.0 m/s

---

Q5.

Fig 1 shows two forces, 12.0 N and 15.0 N, acting on a block of mass  $m = 2.00$  kg. The block slides along a rough horizontal table with coefficient of kinetic friction,  $\mu$  between the block and the table equal to 0.200. Find the acceleration  $a$  of the block.

Fig#



- A)  $2.54 \text{ m/s}^2$
- B)  $5.12 \text{ m/s}^2$
- C)  $7.90 \text{ m/s}^2$
- D)  $9.89 \text{ m/s}^2$
- E)  $1.41 \text{ m/s}^2$

Q6.


The sum of all the external forces on a block is zero. Which one of the following must be true?

- A) The total linear momentum of the block is constant
- B) The acceleration of the block is not zero
- C) The speed of the block is increasing
- D) The block is not in equilibrium
- E) The speed of the block is decreasing

Q7.

A 1000 kg car drives over the top of a circular hill that has a radius of  $R = 50$  m. The speed at the top of the hill is  $v = 20$  m/s. Find the normal force on the car at the top of the hill. (see Fig. 2)

Fig#



- A) 1800 N
- B) 1000 N
- C) 870 N
- D) 1500 N
- E) 2400 N

---

Q8.

A car has a kinetic energy of 25 J. It then makes a U-turn and moves in the opposite direction with twice the original speed. What is the new kinetic energy of the car?

- A) 100J
- B) 50J
- C) -100J
- D) -50J
- E) 25J

---

Q9.

A 60.0 kg student walks up a hill with constant speed reaching a vertical height of 5.00 m above his initial position. How much work does the gravitational force do on him during this walk?

- A) -2940 J
- B) 4950 J
- C) 2500 J
- D) -2500 J
- E) 0 J

---

Q10.

A 3.0 kg box is given an initial speed of 2.2 m/s on a rough horizontal floor. It stops in 2.0 s due to friction between the box and floor. The work done by the frictional force is:

- A) -7.3 J
- B) -9.8 J
- C) -6.5 J
- D) +9.8 J
- E) 0 J

---

Q11.

A 0.40 kg ball moving with a horizontal velocity  $\vec{v}_i = (30 \hat{i})$  m/s hits a vertical wall and bounces back in the opposite direction with velocity  $\vec{v}_f$ . If the impact (collision) of the ball with the wall lasts for 0.10 s and the average force of the wall on the ball is  $-200 \hat{i}$  N, find  $\vec{v}_f$ .

- A)  $-20 \hat{i}$  m/s
- B)  $-30 \hat{i}$  m/s
- C)  $+60 \hat{i}$  m/s
- D)  $+10 \hat{i}$  m/s

E)  $-15 \hat{j}$  m/s

Q12.

Two masses  $m_1 = 3.0$  kg (having velocity  $\vec{v}_1 = 6.0 \hat{i}$  m/s) and  $m_2 = 5.0$  kg (having velocity  $\vec{v}_2 = -6.0 \hat{i}$  m/s) collide and stick together. The final velocity after collision is:

- A)  $-1.5 \hat{i}$  m/s
- B)  $1.5 \hat{i}$  m/s
- C)  $2.0 \hat{i}$  m/s
- D)  $-0.5 \hat{i}$  m/s
- E)  $-2.0 \hat{i}$  m/s

Q13.


A wheel rotates at an angular speed of 600 revolutions per minute around its central axis. It has a rotational kinetic energy of 24000 J about this fixed axis. Calculate the rotational inertia of the wheel about this axis.

- A)  $12 \text{ kg}\cdot\text{m}^2$
- B)  $2.0 \text{ kg}\cdot\text{m}^2$
- C)  $8.5 \text{ kg}\cdot\text{m}^2$
- D)  $14 \text{ kg}\cdot\text{m}^2$
- E)  $10 \text{ kg}\cdot\text{m}^2$

Q14.

A disk of radius  $r = 0.10$  m has a rotational inertia of  $0.020 \text{ kg}\cdot\text{m}^2$  about its axis O (see Fig 3). A string is wound around the disk and pulled with a force of 1.4 N. The angular acceleration of the disk is:

Fig#



- A)  $7.0 \text{ rad/s}^2$
- B)  $3.5 \text{ rad/s}^2$

- C)  $10 \text{ rad/s}^2$
- D)  $14 \text{ rad/s}^2$
- E)  $20 \text{ rad/s}^2$

Q15.


A disk of mass  $5.0 \text{ kg}$  and radius  $0.20 \text{ m}$  rolls smoothly on a horizontal floor. If the kinetic energy of rolling of the disk is  $70 \text{ J}$  at a certain instant, find the speed of the center of mass of the disk. [ $I_{\text{com}}(\text{disk}) = \frac{1}{2} MR^2$ ]

- A)  $4.3 \text{ m/s}$
- B)  $2.5 \text{ m/s}$
- C)  $8.0 \text{ m/s}$
- D)  $40 \text{ m/s}$
- E)  $0 \text{ m/s}$

Q16.

A uniform steel bar of length  $3.0 \text{ m}$  and weight  $20 \text{ N}$  rests on two supports ( $A$  and  $B$ ) at its ends. A block of weight  $W = 30 \text{ N}$  is placed at a distance  $1.0 \text{ m}$  from  $A$  (see Fig. 4). The forces on the supports  $A$  and  $B$  respectively are:

Fig#




- A)  $30 \text{ N}$  and  $20 \text{ N}$
- B)  $25 \text{ N}$  each
- C)  $40 \text{ N}$  and  $10 \text{ N}$
- D)  $35 \text{ N}$  and  $15 \text{ N}$
- E)  $50 \text{ N}$  each

Q17.

Fig. 5 shows a uniform ball of  $600 \text{ N}$  weight suspended by a string  $AB$  and rests against a frictionless vertical wall. The string makes an angle of  $30.0^\circ$  with the wall. The magnitude of the tension in the string is:

Fig#



- A) 693 N
- B) 346 N
- C) 520 N
- D) 300 N
- E) 600 N

Q18.

A horizontal steel rod of length 81 cm and radius 9.5 mm is fixed at one end. It stretches by 0.90 mm when a horizontal force of magnitude  $F$  is applied to its free end. Find the magnitude of  $F$  (Young modulus of steel is  $20 \times 10^{10} \text{ N/m}^2$ ).

- A) 63 kN
- B) 9.8 kN
- C) 0.90 kN
- D) 2.7 kN
- E) 81 kN

Q19.

A spaceship is going from the Earth (mass =  $M_e$ ) to the Moon (mass =  $M_m$ ) along the line joining their centers. At what distance from the centre of the Earth will the net gravitational force on the spaceship be zero? (Assume that  $M_e = 81 M_m$  and the distance from the centre of the Earth to the center of the Moon is  $3.8 \times 10^5 \text{ km}$ ).

- A)  $3.4 \times 10^5 \text{ km}$
- B)  $6.4 \times 10^5 \text{ km}$
- C)  $2.8 \times 10^5 \text{ km}$
- D)  $4.7 \times 10^5 \text{ km}$
- E)  $1.9 \times 10^5 \text{ km}$

Q20.

A 1000 kg satellite is in a circular orbit of radius  $= 2R_e$  about the Earth. How much energy is required to transfer the satellite to an orbit of radius  $= 4R_e$ ? ( $R_e$  = radius of Earth =  $6.37 \times 10^6 \text{ m}$ , mass of the Earth =  $5.98 \times 10^{24} \text{ kg}$ )

- A)  $7.8 \times 10^9$  J.
- B)  $6.1 \times 10^9$  J.
- C)  $4.9 \times 10^8$  J.
- D)  $2.4 \times 10^9$  J.
- E)  $1.7 \times 10^8$  J.

---

Q21.

At what altitude above the Earth's surface would the gravitational acceleration be  $\frac{a_g}{4}$ ?

(where  $a_g$  is the acceleration due to gravitational force at the surface of Earth and  $R_e$  is the radius of the Earth).

- A)  $R_e$
- B)  $2 R_e$
- C)  $R_e/2$
- D)  $R_e/4$
- E)  $3 R_e$

---

Q22.

The gravitational acceleration on the surface of a planet, whose radius is 5000 km, is  $4.0 \text{ m/s}^2$ . The escape speed from the surface of this planet is:

- A) 6.3 km/s
- B) 2.8 km/s
- C) 2.0 km/s
- D) 4.0 km/s
- E) 8.0 km/s

---

Q23.

Water is pumped out of a swimming pool at a speed of 5.0 m/s through a uniform hose of radius 1.0 cm. Find the mass of water pumped out of the pool in one minute. (Density of water =  $1000 \text{ kg/m}^3$ ).


- A) 94 kg
- B) 0.094 kg
- C) 1.6 kg
- D) 19 kg
- E) 5.1 kg

---

Q24.

A large tank open to atmosphere is filled with water. Fig 6 shows this tank with a stream of water flowing through a hole (open to atmosphere) at a depth of 4.00 m. The speed of water,  $v_2$ , leaving the hole is:

Fig#




- A) 8.85 m/s
- B) 4.42 m/s
- C) 2.21 m/s
- D) 17.7 m/s
- E) 35.4 m/s

Q25.

A 10 kg spherical object with a volume of  $0.10 \text{ m}^3$  is held in static equilibrium under water by a cable fixed to the bottom of a water tank. What is the tension  $T$  in the cable? (See Fig. 7)

Fig#



- A) 880 N
- B) 980 N
- C) 1000 N
- D) 1800 N
- E) Zero

Q26.



A plane is at an altitude of 10,000 m where the outside air pressure is 0.25 atm. If the air pressure inside the plane is 1.0 atm, what is the net outward force on  $1\text{ m} \times 2\text{ m}$  door in the wall of the plane?

( $1.0\text{ atm} = 1.01 \times 10^5\text{ Pa}$ ).

- A)  $1.5 \times 10^5\text{ N}$
- B)  $8.5 \times 10^4\text{ N}$
- C) 5.7 N
- D)  $5.9 \times 10^3\text{ N}$
- E)  $1.9 \times 10^{15}\text{ N}$

---

Q27.

A block of mass 20 g is attached to a horizontal spring with spring constant of 25 N/m. The other end of the spring is fixed. The block is pulled a distance 10 cm from its equilibrium position ( $x = 0$ ) on a frictionless horizontal table and released. The frequency of the resulting simple harmonic motion is:

- A) 5.6 Hz
- B) 10 Hz
- C) -10 Hz
- D) 25 Hz
- E) 50 Hz

---

Q28.

A horizontal spring is fixed at one end. A block attached to the other end of the spring undergoes a simple harmonic motion on a frictionless table. Which one of the following statements is correct?

- A) The frequency of the motion is independent of the amplitude of oscillation.
- B) The frequency of the motion is proportional to the amplitude of oscillation.
- C) The acceleration of the block is constant.
- D) The maximum speed of the block is independent of the amplitude.
- E) The maximum acceleration of the block is independent of the amplitude.

---

Q29.

A simple pendulum consists of a mass  $m = 6.00\text{ kg}$  at the end of a light cord of length  $L$ . The angle  $\theta$  between the cord and the vertical is given by  $\theta = 0.08 \cos[(4.43 t + \pi)]$ , where  $t$  is in second and  $\theta$  is in radian. Find the length  $L$ .

- A) 0.50 m
- B) 0.60 m
- C) 0.70 m
- D) 0.80 m
- E) 1.0 m

---

Q30.

A block attached to an ideal horizontal spring undergoes a simple harmonic motion about the equilibrium position ( $x = 0$ ) with an amplitude  $x_m = 10\text{ cm}$ . The mechanical energy of the system is 16 J. What is the kinetic energy of the block when  $x = 5.0\text{ cm}$ ?

- A) 12 J
  - B) 16 J
  - C) 8.0 J
  - D) 4.0 J
  - E) 32 J
-