

PHYS 101 FINAL EXAM-TERM 102


Q1.

A truck moves with a constant speed of 10 m/s in a straight road. It passes point A at time  $t = 0$  and continues towards point B. Ten minutes after the truck passes the point A, a car moving with a constant speed of 15 m/s passes the same point A and continues towards B along the same straight road. The car will catch up with the truck at time  $t$  equals to:

- A) 30 minutes
- B) 60 minutes
- C) 3 minutes
- D) 10 minutes
- E) 15 minutes

Q2.

**Figure 1** shows the position-time graph for two objects, A and B, moving along a straight line. Which one of the following statements is TRUE?



- A) The speed of B is always greater than the speed of A.
- B) The two objects have the same speed at  $t = 4$  s.
- C) Object B is always ahead of object A.
- D) Object A is always ahead of object B.
- E) The speed of A is always greater than the speed of B.

Q3.

Consider two vectors  $\vec{v} = 3.0 \hat{i} + 3.0 \hat{j}$  and  $\vec{w} = \cos \theta \hat{i} + \sin \theta \hat{j}$ , where  $\theta$  is measured counter clockwise with respect to the positive  $x$ -axis. For what value of  $\theta$  (in degrees) is  $\vec{v} \times \vec{w} = 0$ ?

- A) 45
- B) 135
- C) 90
- D) 105
- E) 0

Q4.


A 2-kg object is initially at rest. At time  $t = 0$ , a force  $\vec{F}_1 = (2\hat{i} + 2\hat{j})N$ , is applied to the object. At time  $t = 1$  s, an additional force  $\vec{F}_2 = (-2\hat{i} - 2\hat{j})N$  is applied to the object. Find the velocity of the object at  $t = 2$  s.

- A)  $(\hat{i} + \hat{j})m/s$
- B)  $(-\hat{i} - \hat{j})m/s$
- C)  $(2\hat{i} + 2\hat{j})m/s$
- D)  $(-2\hat{i} - 2\hat{j})m/s$
- E) 0

Q5.

A force  $\vec{F}$  is applied to a block of mass equal to 3.00 kg resting on a rough horizontal surface. The force makes an angle of  $37.0^\circ$  with the horizontal as shown in **Figure 2**. The coefficient of static friction between the block and the surface is 0.450. If the block is just about to slide, calculate the magnitude of the force  $\vec{F}$ .

Fig#




- A) 12.4 N
- B) 19.6 N
- C) 16.5 N
- D) 10.7 N
- E) 20.6 N

Q6.

The system shown in **Figure 3** is released from rest and is moving with an acceleration of  $4.00 \text{ m/s}^2$ . Find the magnitude of the tension  $T$  shown in the figure. (Assume that the pulley and the cords are massless).

Fig#



- A)  $T = 11.6 \text{ N}$
- B)  $T = 6.96 \text{ N}$
- C)  $T = 15.4 \text{ N}$
- D)  $T = 10.0 \text{ N}$
- E)  $T = 4.80 \text{ N}$

Q7.

If the weight of an object on the Moon is one-sixth of its weight on Earth, the ratio of its kinetic energy when it is moving with speed  $V$  on Earth to its kinetic energy when it is moving with the same speed  $V$  on the Moon is:

- A) 1.0
- B) 6.0
- C) 2.6
- D) 3.1
- E) 1.6

Q8.

A block is released from rest at the top of an inclined plane making an angle of  $30.0^\circ$  with the horizontal. The coefficient of kinetic friction between the block and the inclined plane is 0.300. What is the speed of the block after it has traveled a distance of 1.00 m downwards along the inclined plane?

- A) 2.17 m/s
- B) 3.58 m/s
- C) 4.30 m/s
- D) 5.57 m/s
- E) 7.33 m/s

Q9.

A  $1.00 \times 10^3 \text{ kg}$  car is traveling at 20.0 m/s toward the north. During a collision, the car receives an impulse of magnitude  $1.00 \times 10^4 \text{ N}\cdot\text{s}$  toward the south. What is the velocity of the car immediately after the collision?

- A) 10.0 m/s, north
- B) 30.0 m/s, north

- C) 20.0 m/s, north
- D) 10.0 m/s, south
- E) 20.0 m/s, south

Q10.


Two blocks approach each other at right angles on a frictionless surface. Block A has a mass of 45.1 kg and travels in the  $+x$  direction at 3.20 m/s. Block B has a mass of 85.8 kg and is moving in the  $+y$  direction at 2.08 m/s. They collide and stick together. Find the final velocity of the two blocks.

- A)  $(1.10 \hat{i} + 1.36 \hat{j})$  m/s
- B)  $(2.30 \hat{i} + 3.36 \hat{j})$  m/s
- C)  $(3.45 \hat{i} + 2.56 \hat{j})$  m/s
- D)  $(5.20 \hat{i} + 6.37 \hat{j})$  m/s
- E)  $(4.50 \hat{i} + 4.76 \hat{j})$  m/s

Q11.

As shown in **Figure 4**, a 45-N force is applied to one end of a massless string which is wrapped around a pulley that has a radius of 1.5 m and a moment of inertia of  $2.25 \text{ kg}\cdot\text{m}^2$ . Through what angle will the pulley rotate in 3.0 s if it was initially at rest?

Fig#




- A) 135 rad
- B) 90.0 rad
- C) 451 rad
- D) 270 rad
- E) 225 rad

Q12.

**Figure 5** shows a uniform horizontal beam of mass  $M = 4.00 \text{ kg}$  and length  $L = 4.00 \text{ m}$  being acted upon by four forces of magnitudes  $F_1 = 10.0 \text{ N}$ ,  $F_2 = 20.0 \text{ N}$ ,  $F_3 = 30.0 \text{ N}$  and  $F_4 = 10.0 \text{ N}$  and in the directions as indicated. Find the net torque about point O at the center of the beam.

Fig#




- A) 10.0 N.m, counter clockwise
- B) 10.0 N.m, clockwise
- C) 100 N.m, counter clockwise
- D) 100 N.m, clockwise
- E) 140 N.m, counter clockwise

Q13.

As shown in **Figure 6**, a uniform beam of length 4.20 m is suspended by a cable from its center point O. A 65.0-kg man stands at one end of the beam. Where should a 190-kg block be placed on the beam so that the beam is in static equilibrium (Distances are measured from the center point O of the beam)?

Fig#



- A) 0.718 m
- B) 1.44 m
- C) 2.35 m
- D) 0.543 m
- E) 2.10 m

Q14.


What increase in pressure is necessary to decrease the volume of a sphere by 0.150 % (Take the bulk modulus of the sphere  $B = 2.80 \times 10^{10} \text{ N/m}^2$ )?

- A)  $4.20 \times 10^7 \text{ N/m}^2$
- B)  $1.40 \times 10^7 \text{ N/m}^2$
- C)  $3.56 \times 10^6 \text{ N/m}^2$
- D)  $2.80 \times 10^7 \text{ N/m}^2$
- E)  $1.01 \times 10^5 \text{ N/m}^2$

Q15.

Three parallel forces of magnitudes 10.0 N, 20.0 N, and 20.0 N, respectively, act on a body (**Figure 7**). The perpendicular distances from a given point O to their lines of action are shown. The single force which can replace these forces is:

Fig#




- A) 50.0 N, 10.0 cm to the right of point O.
- B) 50.0 N, 20.0 cm to the right of point O.
- C) 30.0 N, 17.5 cm to the right of point O.
- D) 50.0 N, 17.5 cm to the right of point O.
- E) 50.0 N, acting through the given point O.

Q16.

Five masses are put together as shown in **Figure 8**. What is the net force on the 1.0-kg mass placed in the center of the circle?  $G$  is the gravitational constant.

Fig#



- A)  $G/a^2 (+\hat{j})$
- B)  $G/a^2 (-\hat{j})$
- C) 0
- D)  $3G/a^2 (\hat{i} + \hat{j})$
- E)  $4G/a^2 (-\hat{j})$

Q17.

If, instead of being distributed over the volume of the Earth, the mass of the Earth is distributed inside a thin shell, what would be the radial dependence of the gravitational force on an object outside the Earth? Take  $r$  to be the distance to the object from the center of the Earth.

- A)  $1/r^2$
- B)  $1/r$
- C)  $1/r^3$
- D)  $1/\sqrt{r}$
- E) None of the others

Q18.

If we assume that a black hole is a planet where the escape velocity is equal to the speed of light ( $3.00 \times 10^8$  m/s), find the radius of a black hole with a mass equal to that of Earth.

- A)  $8.86 \times 10^{-3}$  m
- B)  $8.85 \times 10^{+3}$  m
- C)  $6.38 \times 10^{+3}$  m
- D)  $6.38 \times 10^{-3}$  m
- E)  $3.00 \times 10^{+8}$  m

Sec# 13-6

Grade# 60

Q19.

The law of areas due to Kepler is equivalent to the law of

- A) Conservation of angular momentum.
- B) Conservation of mass.
- C) Conservation of energy.
- D) Conservation of linear momentum.
- E) None of the others.

Q20.

What speed on the surface of Earth should be given to a satellite to put it in an orbit of radius  $R = 3R_E$  around the Earth (where  $R_E$  is the radius of Earth)?


- A)  $\sqrt{\frac{10 G M_E}{6 R_E}}$
- B)  $\sqrt{\frac{5 G M_E}{6 R_E}}$
- C)  $\sqrt{\frac{8 G M_E}{6 R_E}}$
- D)  $\sqrt{\frac{G M_E}{R_E}}$

E)  $\sqrt{\frac{7 G M_E}{6 R_E}}$

Q21.

In the hydraulic lift of **Figure 9**, a large piston of diameter  $d_1 = 120$  cm supports a car of mass  $3.20 \times 10^3$  kg. What is the magnitude of the vertically downward force  $F_1$  that must be applied to the smaller piston of diameter  $d_2 = 15.0$  cm to balance the car?

Fig#




- A)  $4.90 \times 10^2$  N
- B)  $3.92 \times 10^3$  N
- C)  $1.50 \times 10^3$  N
- D)  $2.00 \times 10^2$  N
- E)  $2.50 \times 10^4$  N

Q22.

A U-shaped tube open at both ends contains water and a quantity of oil occupying a 2.0 cm length of the tube, as shown in **Figure 10**. If the density of oil is 82% of the density of water, what is the height difference  $h$ ?

Fig#





- A) 0.36 cm
- B) 1.2 cm
- C) 0.43 cm
- D) 0.75 cm
- E) 0.82 cm

Q23.

The average density of a typical iceberg is 0.86 that of sea water. What fraction of the volume of the iceberg is outside the water?

- A) 0.14
- B) 0.86
- C) 0.50
- D) 0.45
- E) 0.75

Q24.

Water flows through a horizontal pipe of varying cross-section. The pressure is  $1.5 \times 10^4$  Pa at a point where the speed is 2.0 m/s and the area of cross section is A. Find the speed and pressure at a point where the area is A/2.


- A) 4.0 m/s and  $0.90 \times 10^4$  Pa
- B) 4.0 m/s and  $0.75 \times 10^4$  Pa
- C) 8.0 m/s and  $0.90 \times 10^4$  Pa
- D) 8.0 m/s and  $1.5 \times 10^4$  Pa
- E) 2.0 m/s and  $1.8 \times 10^4$  Pa

Sec# 14-9  
Grade# 43

Q25.

A large tank is filled with water. A tightly fitting piston rests on top of the water (**Figure 11**). The combined pressure from the piston and atmosphere on the top surface of water is  $1.02 \times 10^5$  Pa. A very small circular hole is opened at a depth of 60.0 cm below the initial water level of the tank. What is the initial speed of water coming out of the hole?

Fig#



- A) 3.71 m/s
- B) 5.43 m/s
- C) 9.80 m/s
- D) 4.93 m/s
- E) 1.60 m/s


Q26.

If the amplitude of oscillation of an object in simple harmonic motion is increased, then

- A) the total mechanical energy of the object will increase
- B) the period of oscillations of the object will increase
- C) the frequency of oscillations of the object will increase
- D) the frequency of oscillations of the object will decrease
- E) the maximum kinetic energy of the object will decrease

Q27.

A solid circular disk is oscillating with a period  $T$  in a vertical plane about pivot point  $P$  as shown in **Figure 12**. If the disk is made four times heavier but still having the same radius, what will be its period of oscillation?



- A)  $T$
- B)  $2T$
- C)  $T/2$
- D)  $T/4$
- E)  $4T$

Q28.

The maximum speed of a 3.00-kg object executing simple harmonic motion is 6.00 m/s. The maximum acceleration of the object is  $5.00 \text{ m/s}^2$ . What is its period of oscillations?

- A) 7.54 s
- B) 2.50 s
- C) 1.20 s
- D) 0.833 s
- E) 0.278 s

Q29.

An object executes simple harmonic motion with an amplitude of 1.2 cm and a time period of 0.10 s. What is the total distance traveled by the object in 1.9 s?

- A) 91 cm
- B) 27 cm
- C) 40 cm
- D) 11 cm
- E) 70 cm

Q30.

A simple pendulum of length  $L_1$  has time period  $T_1$ . A second simple pendulum of length  $L_2$  has time period  $T_2$ . If  $T_2 = 2 T_1$ , find the ratio  $L_1/L_2$ .

- A)  $1/4$
- B)  $1/2$
- C) 4
- D) 2
- E) 1