

Q1.

A ball is thrown from the ground vertically upward and reaches a maximum height of 40 m. Upon descending, the ball hits the ground and rebounds with half its initial velocity. Find the height to which it reaches after rebounding.

- A) 10 m
- B) 20 m
- C) 15 m
- D) 8.8 m
- E) 25 m

Q2.

You are given vectors $\mathbf{A} = 5.00 \mathbf{i} - 6.00 \mathbf{j}$ and $\mathbf{B} = -3.00 \mathbf{i} + 7.00 \mathbf{j}$. A third vector \mathbf{C} lies in the xy-plane. Vector \mathbf{C} is perpendicular to vector \mathbf{A} , and the scalar product of \mathbf{C} with \mathbf{B} is 15.0. Find the vector \mathbf{C} .

- A) $5.29 \mathbf{i} + 4.41 \mathbf{j}$
- B) $7.54 \mathbf{i} + 3.42 \mathbf{j}$
- C) $6.32 \mathbf{i} + 2.53 \mathbf{j}$
- D) $8.37 \mathbf{i} + 4.24 \mathbf{j}$
- E) $7.21 \mathbf{i} + 2.23 \mathbf{j}$

Q3.


An object is moving on a horizontal circular path of radius 1.5 meters at a constant speed. The time required for one revolution is 3.2 s. The acceleration of the object is:

- A) 5.8 m/s^2
- B) 2.6 m/s^2
- C) 7.7 m/s^2
- D) 1.4 m/s^2
- E) zero

Q4.

A projectile is fired with initial velocity \mathbf{v}_0 and angle $\theta_0 = 60^\circ$ from the top of a building (**Figure 1**) and is observed to reach a maximum height of 19.6 m. It later hits the ground at a horizontal distance of 50.0 m from the base of the building. Find the time of flight of the projectile. (Neglect air friction)

Fig#




- A) 4.42 s
- B) 5.00 s
- C) 9.80 s
- D) 3.32 s
- E) 2.50 s

Q5.

As shown in **Figure 2**, a block with mass $M = 3.00$ kg is lying on a smooth surface and is attached to another block of mass $m = 2.00$ kg by means of a light, inextensible string which passes over a massless pulley. What force F acting on the block M at angle $\theta = 60^\circ$ above the horizontal will hold both objects at rest?

Fig#



- A) 39.2 N
- B) 29.4 N
- C) 19.6 N
- D) 49.0 N
- E) 9.80 N

Q6.

A car is moving on a flat horizontal circular track of radius $R = 25.0$ m. The coefficient of static friction between the car wheels and the track is $\mu_s = 0.350$. What is the speed at which the car starts sliding outside the track?


- A) 9.26 m/s

- B) 13.0 m/s
- C) 14.5 m/s
- D) 11.1 m/s
- E) 5.44 m/s

Q7.

A block of mass $m = 5.00$ kg slides on a horizontal rough surface under the action of a steady force F applied to the block at a constant angle of $\theta = 45^\circ$ (**Figure 3**). The coefficient of kinetic friction between the block and the surface is 0.400. Find the maximum value of the force F for which the block will move only horizontally.

Fig#



- A) 69.3 N
- B) 19.6 N
- C) 50.0 N
- D) 98.0 N
- E) 150 N

Q8.


A 100 kg parachute falls at a constant speed of 0.750 m/s. At what rate is energy being lost?

- A) 735 W
- B) 75.0 W
- C) 56.3 W
- D) 28.0 W
- E) 147 W

Q9.

A mass $m = 1.00$ kg is released from rest at point P ($h = 6.00$ m). It slides along the smooth track and reaches point O on the circular part of the track ($R = 1.00$ m) shown in **Figure 4**. What force does the track exert on the mass at point O?

Fig#



- A) 68.6 N
- B) 88.2 N
- C) 98.0 N
- D) 19.6 N
- E) 49.0 N

Q10.

A stationary object of mass $m = 24.0$ kg explodes into two pieces of masses 14.0 kg and 10.0 kg. The velocity of the 10.0 kg mass is 6.00 m/s in the positive x -direction. The change in the kinetic energy of the object is:

- A) 309 J
- B) 511 J
- C) 240 J
- D) 180 J
- E) 160 J

Q11.

If the kinetic energy of a body is increased by a factor of 9, then the momentum of the body will increase by a factor of:

- A) 3
- B) 1
- C) 9
- D) 4
- E) 2

Q12.

A solid spherical ball of mass $M = 0.40$ kg and radius $R = 5.0$ cm is rotating about its fixed central axis with angular speed of 4.0 rad/s. It was brought to a stop in 6.0 s. The work done to stop the ball is:

- A) -3.2×10^{-3} J
- B) -4.8×10^{-3} J
- C) -5.4×10^{-3} J
- D) -1.8×10^{-3} J

E) $-2.8 \times 10^{-3} \text{ J}$

Q13.

An electric fan is turned off, and its angular velocity decreases uniformly from 500 rev/min to 250 rev/min in 4.00 s. Find the number of revolutions made by the motor in the 4.00 s interval.

- A) 25.0
- B) 10.0
- C) 15.0
- D) 20.0
- E) 30.0

Q14.

A 0.20 kg stone attached to a string is rotating with a constant angular speed of 3.0 rev/s in a horizontal circle of radius 0.75 m. The magnitude of the angular momentum of the stone relative to the center of the circle is:

- A) $2.1 \text{ kg}\cdot\text{m}^2/\text{s}$
- B) $4.2 \text{ kg}\cdot\text{m}^2/\text{s}$
- C) $3.2 \text{ kg}\cdot\text{m}^2/\text{s}$
- D) $0.44 \text{ kg}\cdot\text{m}^2/\text{s}$
- E) $1.6 \text{ kg}\cdot\text{m}^2/\text{s}$

Q15.

A hoop rolls down an inclined plane. The ratio of its rotational kinetic energy to its total kinetic energy is:

- A) 1/2
- B) 2/3
- C) 1/3
- D) 1/4
- E) 2

Q16.


A small sphere is made of a material with a bulk modulus of $1.90 \times 10^9 \text{ Pa}$. The volume of the sphere shrinks by 0.20 % when submerged in a fluid at a depth of 400 m. What is the density of this fluid? Assume the pressure on the sphere at this depth is the same from all directions.

- A) 970 kg/m^3
- B) 1200 kg/m^3
- C) 990 kg/m^3
- D) 1000 kg/m^3
- E) 1100 kg/m^3

Q17.

Figure 5 shows a uniform rod (mass $M = 5.0$ kg, length $L = 1.2$ m) dangling over a frictionless edge at point P, but secured by a horizontal rope with tension $T = 5.0$ N. If the angle $\theta = 30^\circ$, then what is the distance d needed to keep the rod in equilibrium.

Fig#




- A) 3.3 cm
- B) 2.6 cm
- C) 4.7 cm
- D) 2.0 cm
- E) 5.8 cm

Q18.

The uniform sphere in **Figure 6** has a mass $m = 2.0$ kg and is held in place by a massless rope of length $L = 20$ cm, touching a rough wall. Find the force of friction between the sphere and the wall if the tension in the rope is 10 N, and the angle $\theta = 30^\circ$.

Fig#



- A) 11 N
- B) 7.0 N
- C) 8.0 N
- D) 18 N
- E) 4.5 N

Q19.

Planet Pluto has a radius 20% of the earth radius and a mass only 0.2% that of earth. If an astronaut can jump 0.5 m high on earth, then how high can he jump on Pluto? (assume the astronaut jumps on both planets with the same velocity)

- A) 10 m
- B) 20 m
- C) 0.5 m
- D) 5.0 m
- E) 0.05

Q20.


A satellite orbits a planet of unknown mass in a circle of radius 2.0×10^7 m. The magnitude of the gravitational force on the satellite is 80 N. What is the kinetic energy of the satellite in this orbit?

- A) 80×10^7 J
- B) 2.5×10^5 J
- C) 40×10^7 J
- D) 1.6×10^7 J
- E) 32×10^7 J

Q21.

Figure 7 shows two particles of masses, m and $2m$ fixed in their positions. A particle of mass m is to be brought from an infinite distance to one of the three locations, a , b and c . Rank these three locations according to the magnitude of the net work done by the gravitational force on this particle due to the fixed particles, greatest first.

Fig#



- A) b, a, c
- B) b , then a and c tie
- C) a, c, b
- D) c, a, b
- E) all tie

Q22.

Which one of the following statements concerning Kepler's laws is FALSE?

- A) Satellites in the same orbit around the earth but with different masses will have different periods.

- B) Satellites with the same masses but in different orbits having different radii around the earth will have different periods.
- C) The angular momentum is conserved for planets rotating about the sun.
- D) The planets move faster when they are close to the sun.
- E) Planets in their orbits sweep equal areas in equal times.

Q23.


The volume flow rate of water through a horizontal pipe is $2.0 \text{ m}^3/\text{min}$. Calculate the speed of flow at a point where the radius of the pipe is 10 cm.

- A) 1.1 m/s
- B) 3.2 m/s
- C) 0.55 m/s
- D) 2.5 m/s
- E) 4.0 m/s

Q24.

A uniform U-tube is partially filled with water. Oil, of density 0.75 g/cm^3 , is poured into the right arm as shown in **Figure 8**. The length of the oil column (h) is then:

Fig#




- A) 4.0 cm
- B) 8.0 cm
- C) 6.0 cm
- D) 2.0 cm
- E) 10 cm

Q25.

A disk made of lead (diameter = 5.0 cm, height = 3.0 cm, density = $11.3 \times 10^3 \text{ kg/m}^3$) floats in a container of mercury (density = $13.6 \times 10^3 \text{ kg/m}^3$). What is the depth x (see **Figure 9**) by which the disk sinks in mercury.

Fig#




- A) 2.5 cm
- B) 1.3 cm
- C) 2.0 cm
- D) 2.8 cm
- E) 1.7 cm

Q26.

The edge length of the cube in **Figure 10** is 10 cm and its mass is 2.0 kg. It hangs from a spring and is fully submerged in water. If the spring constant is 98 N/m, by how much does the spring stretch from its equilibrium length.

Fig#




- A) 10 cm
- B) 20 cm
- C) 15 cm
- D) 5.0 cm
- E) 25 cm

Q27.

A simple pendulum of length 12 cm is to be replaced by a hoop in one of the old O'clocks. See **Figure 11**. What should be the radius of the hoop needed to produce the same period as that of the pendulum, while oscillating about point O?

Fig#




- A) 6.0 cm
- B) 4.0 cm
- C) 3.0 cm
- D) 12 cm
- E) 24 cm

Q28.

The velocity versus time plot for a block-spring system performing a simple harmonic motion is shown in **Figure 12**. The horizontal scale is set by $t_s = 0.2$ s. Find the acceleration of the system at $t = 0.1$ s.

Fig#



- A) 200 m/s^2
- B) 100 m/s^2
- C) 80 m/s^2
- D) 10 m/s^2
- E) 60 m/s^2

Q29.

A block-spring system is in simple harmonic motion and its displacement as a function of time is given by the equation:

$$x = (5.0 \text{ m}) \cos[(\pi/3 \text{ rad/s})t - \pi/4 \text{ rad}],$$

The mass of the block is 3.0 kg. Find the speed of the block when the kinetic energy is one-fourth the total energy.

- A) 2.6 m/s
- B) 4.7 m/s
- C) 3.3 m/s
- D) 5.0 m/s

E) 1.5 m/s

Q30.

A mass $m_1 = 1.0$ kg is connected to a spring (with spring constant equal to k) and oscillates on a horizontal frictionless table with a period of 1.0 s. When m_1 is replaced with another unknown mass m_2 , the period changes to 2.0 s. Find the value of m_2 .

- A) 4.0 kg
 - B) 3.0 kg
 - C) 2.0 kg
 - D) 0.50 kg
 - E) 1.0 kg
-