

Q1.

A car accelerates at 2.0 m/s^2 along a straight road. It passes two marks that are 30 m apart at times $t = 4.0 \text{ s}$ and $t = 5.0 \text{ s}$. Find the car's velocity at $t = 0$.

- A) 21 m/s
- B) 34 m/s
- C) 16 m/s
- D) 11 m/s
- E) 48 m/s

Q2.

Two vectors are given by $\vec{A} = 1.00\hat{i} + 2.00\hat{j}$ and $\vec{B} = 1.00\hat{i} + 3.00\hat{j}$. Find the angle that the vector $\vec{A} - 2\vec{B}$ makes with the positive y -axis.

- A) 166°
- B) 100°
- C) 133°
- D) 111°
- E) 173°

Q3.

A projectile's launch speed is 4 times its speed at maximum height. Find the launch angle from the horizontal.

- A) 75.5°
- B) 70.6°
- C) 45.3°
- D) 32.0°
- E) 49.2°

Q4.

A particle moves at constant speed in a circular path. The instantaneous velocity and instantaneous acceleration vectors are both:

- A) Perpendicular to each other
 - B) Perpendicular to the circular path
 - C) tangent to the circular path
 - D) Opposite to each other
 - E) Parallel to each other
-

Q5.

An elevator initially moving upward is slowing down at a rate of 1.50 m/s^2 . If the tension in the cable is $3.20 \times 10^3 \text{ N}$ then find the weight of the elevator.

- A) $3.78 \times 10^3 \text{ N}$
- B) $1.53 \times 10^4 \text{ N}$
- C) $5.20 \times 10^5 \text{ N}$
- D) $1.72 \times 10^3 \text{ N}$
- E) $5.92 \times 10^3 \text{ N}$

Q6.

A 12 N horizontal force is applied to a 4.1 kg block initially at rest on a rough horizontal surface. If the coefficients of friction are $\mu_s = 0.5$ and $\mu_k = 0.4$. Find the magnitude of the frictional force on the block.

- A) 12 N
- B) 16 N
- C) 10 N
- D) 20 N
- E) 8.0 N

Q7.

A single force F acts on a block of mass $m = 3.0 \text{ kg}$ from $t = 0 \text{ s}$ to $t = 4.0 \text{ s}$. If the position of the block is given by $x = t^3 - 5.2t$ then find the work done on the block by F .

- A) $2.7 \times 10^3 \text{ J}$
- B) $5.4 \times 10^3 \text{ J}$
- C) $4.2 \times 10^3 \text{ J}$
- D) $6.7 \times 10^3 \text{ J}$
- E) $1.2 \times 10^3 \text{ J}$

Q8.

The Rotational inertia of an object does not depend upon:

- A) Its angular velocity.
 - B) Its mass.
 - C) Its size and shape.
 - D) The location of the axis of rotation.
 - E) The distribution of its mass.
-

Q9.

A uniform meter stick pivoted at 10.0 cm mark is oscillating. Find the period of oscillation.

- A) 1.57 s
- B) 2.32 s
- C) 3.60 s
- D) 4.15 s
- E) 3.43 s

Q10.

A thin uniform rod of length 1.5 m and mass 0.50 kg is suspended freely from one end. It is pulled to one side and then allowed to swing like a pendulum, passing through its lowest position with angular speed 5.0 rad/s. Neglecting friction and air resistance, find the rod's kinetic energy at its lowest position.

- A) 4.7 J
- B) 1.2 J
- C) 9.4 J
- D) 0.90 J
- E) 7.8 J

Q11.


A force of magnitude 10.0 N acts on a rigid body. The force lies in the xy plane. Its line of action passes through the point (0.500, 0.00) and makes an angle of 30.0° with the positive x -axis. Find the torque of the force about the point (-0.300, 0.00).

- A) $+4.00 \hat{k}$ (N.m)
 - B) $-4.00 \hat{k}$ (N.m)
 - C) $+1.00 \hat{k}$ (N.m)
 - D) $-1.00 \hat{k}$ (N.m)
 - E) $+6.93 \hat{i}$ (N.m)
-

Q12.

A horizontal uniform beam of weight 1000 N is supported by a hinge at one end and by a cable at the other end, as shown in **Figure 1**. Find the magnitude of the force exerted on the beam by the hinge.

Fig#




- A) 1000 N
- B) 1200 N
- C) 780.0 N
- D) 1500 N
- E) 892.0 N

Q13.

Four particles, each with mass m , are arranged symmetrically about the origin on the x axis, as shown in **Figure 2**. A fifth particle, with mass M , is on the y axis. The direction of the gravitational force on M is:

Fig#



- A) Along the negative y axis
- B) Along the positive y axis
- C) Along the negative x axis
- D) Along the positive x axis
- E) Along the negative z axis

Q14.

A uniform solid sphere has a mass of 1.5×10^4 kg and a radius of 1.0 m. Find the magnitude of the gravitational force due to the sphere on a particle of mass $m = 1.0$ kg located at a distance of 0.75 m from the center of the sphere.

- A) 7.5×10^{-7} N
- B) 1.9×10^{-7} N
- C) 3.6×10^{-7} N
- D) 9.9×10^{-7} N
- E) 0

Q15.

A planet has a mass of about 0.0558 times the mass of Earth and a diameter of about 0.381 times the diameter of Earth. The acceleration of a body falling near the surface of this planet is: (take acceleration due to gravity on earth to be 9.8 m/s^2)

- A) 3.77 m/s^2
- B) 1.50 m/s^2
- C) 5.95 m/s^2
- D) 9.80 m/s^2
- E) 2.42 m/s^2

Q16.


Neglecting air resistance, a 1.0 kg projectile has an escape speed of about 11 km/s at the surface of Earth. Find the corresponding escape speed for a 2.0 kg projectile.

- A) 11 km/s
 - B) 7.2 km/s
 - C) 15 km/s
 - D) 5.5 km/s
 - E) 22 km/s
-

Q17.

Two satellites *A* and *B* of same mass of 130 kg are shown in **Figure 3**, and move in the same circular orbit of radius $r = 7.77 \times 10^6$ m around earth but of opposite senses of rotation and therefore they are expected to collide. If the collision is completely inelastic, find the total mechanical energy immediately after collision.

Fig#



- A) -1.33×10^{10} J
- B) -3.39×10^{10} J
- C) -1.98×10^8 J
- D) -2.93×10^{10} J
- E) 0

Q18.


At a fixed depth within a fluid at rest, the pressure pushing upward is:

- A) Equal to pressure pushing downward.
 - B) Zero, because pressure only pushes equal in all horizontal direction.
 - C) Zero, because the fluid above does not support the weight of the fluid below.
 - D) Greater than the pressure pushing downward.
 - E) Less than the pressure pushing downward.
-

Q19.

A 5.0 kg rock whose density is 4800 kg/m^3 is suspended by a string such that half of the rock's volume is under water (see **Figure 4**). Find the tension in the string.

Fig#




- A) 44 N
- B) 73 N
- C) 32 N
- D) 68 N
- E) 21 N

Q20.

Figure 5 shows volume flow rates (in cm^3/s) of a fluid from all but one tube. Assuming steady flow of the fluid, find the volume flow rate through the **X** tube and its direction.

Fig#




- A) 1 cm^3/s flowing out
- B) 7 cm^3/s flowing in
- C) 5 cm^3/s flowing out
- D) 3 cm^3/s flowing in
- E) 4 cm^3/s flowing out

Q21.

A 4.0 mm diameter hole is 1.0 m below the surface of a large tank of water as shown in **Figure 6**. Find water volume flow rate through the hole.

Fig#



- A) $5.6 \times 10^{-5} \text{ m}^3/\text{s}$
- B) $3.1 \times 10^{-5} \text{ m}^3/\text{s}$
- C) $7.8 \times 10^{-5} \text{ m}^3/\text{s}$
- D) $1.5 \times 10^{-6} \text{ m}^3/\text{s}$
- E) $4.7 \times 10^{-6} \text{ m}^3/\text{s}$

Q22.


For an object undergoing a simple harmonic motion. Only one statement is correct

- A) The object has varying acceleration.
- B) The object has varying amplitude.
- C) The object has varying period.
- D) The object has varying frequency.
- E) The object has varying total mechanical energy.

Q23.

As shown in **Figure 7**, two identical springs of spring constant $7.00 \times 10^3 \text{ N/m}$ are attached to a block that is sitting on a frictionless floor. If the frequency of oscillation is 30.0 Hz, find the mass of the block.

Fig#



- A) 0.394 kg
- B) 0.126 kg
- C) 0.328 kg
- D) 0.200 kg
- E) 0.175 kg

Q24.

A 3.000 kg block, attached to a spring, executes simple harmonic motion. The position of the block is given as: $x = 2.000\cos(50.00t)$ where x is in meters and t is in seconds. Find the spring constant of the spring:

- A) 7500 N/m
- B) 6800 N/m
- C) 9000 N/m
- D) 2560 N/m
- E) 4700 N/m

Q25.

A 50 kg boy stands on frictionless level ice floor. He kicks a 0.10 kg stone lying near his feet if the velocity of the stone is $(1.1 \text{ m/s})\hat{i}$, find the velocity of the boy just after kicking the stone.

- A) $(-2.2 \times 10^{-3} \text{ m/s})\hat{i}$
 - B) $(2.0 \times 10^{-3} \text{ m/s})\hat{i}$
 - C) $(1.1 \times 10^{-3} \text{ m/s})\hat{i}$
 - D) $(-1.2 \times 10^{-3} \text{ m/s})\hat{i}$
 - E) 0
-