

Q1.

A car travels with constant acceleration along a straight road. How much time does the car take to increase its speed from 30 m/s to 50 m/s in a distance of 180 m?

A) 4.5 s

B) 6.0 s

C) 3.6 s

D) 7.2 s

E) 9.0 s

Q2.

Consider three vectors \vec{A} , \vec{B} , \vec{C} such that $\vec{C} = \vec{A} + \vec{B}$. Which of the following operations will not change the magnitude of \vec{C} ?

A) Rotate \vec{A} and \vec{B} each through the same angle about the same axis

B) Multiply \vec{A} by 2 and divide \vec{B} by 2

C) Divide \vec{A} by 2 and divide \vec{B} by 2

D) Replace \vec{B} by $-\vec{B}$

E) Replace \vec{A} by $-\vec{A}$

Q3.

A ball is thrown horizontally from the top of a building that is 20.0 m high with a speed of 30.0 m/s. What is the speed of the ball when it hits the ground?

A) 35.9 m/s

B) 9.80 m/s

C) 37.3 m/s

D) 30.0 m/s

E) 38.6 m/s

Q4.

Two cars (A and B) are travelling due east on a highway. Their speeds relative to the ground are: $v_A = 30$ km/h and $v_B = 50$ km/h. What is the velocity of A relative to B?

A) 20 km/h due west

B) 20 km/h due east


C) 80 km/h due west

D) 80 km/h due east

E) 40 km/h due east

Q5.

A force F pushes on three blocks on a frictionless surface, as shown in **Figure 1**. What is the magnitude of the force of block B on block A?



A) 35 N

B) 14 N

C) 63 N

D) 21 N

E) 33 N

Q6.

A box is sliding down an incline that is 35° above the horizontal. If the coefficient of kinetic friction between the block and the surface is 0.40, the magnitude of its acceleration is

A) 2.4 m/s^2

B) 5.6 m/s^2

C) 8.8 m/s^2

D) 1.3 m/s^2

E) zero

Q7.

A 2000-kg elevator starts from rest and accelerates upward at 3.20 m/s^2 . How much power is delivered by the tension in the cable pulling the elevator at the instant when it has a speed of 6.00 m/s ?

A) 156 kW

B) 221 kW


C) 64.0 kW

D) 193 kW

E) 48.0 kW

Q8.


A 3.0-kg block starts from rest and slides down a frictionless 30° incline, where it collides with a massless spring of force constant 400 N/m, as shown in **Figure 2**. The block slides a total distance of 0.65 m on the incline until it is stopped by the spring. By how much is the spring compressed?



- A) 0.22 m
- B) 0.37 m
- C) 0.13 m
- D) 0.48 m
- E) 0.31 m

Q9.


In **Figure 3**, a block is released from rest at point A and comes to rest at point C. The track from A to B is frictionless, while the track from B to C is rough. What is the coefficient of kinetic friction between the block and track BC?



- A) 0.53
- B) 0.34
- C) 0.21
- D) 0.72
- E) 0.43

Q10.

A 4.0-kg object moving with speed 30 m/s strikes a surface at angle $\theta = 45^\circ$ and rebounds at the same angle with the same speed (see **Figure 4**). The impulse on the object is



- A) 170 N.s along the + y-axis
- B) 170 N.s along the - y-axis
- C) 340 N.s along the + y-axis
- D) 340 N.s along the - y-axis
- E) 140 N.s at 45° relative to the + x-axis

Q11.

A 4.0-kg object has a velocity of 4.0 m/s in the positive x direction when it explodes into two objects each with a mass of 2.0 kg. After the explosion, one of the objects has a velocity of 3.0 m/s at an angle of 60° measured counterclockwise from the positive x axis. What is the speed of the other object after the explosion?

- A) 7.0 m/s
- B) 8.9 m/s
- C) 7.9 m/s
- D) 6.1 m/s
- E) 6.7 m/s

Q12.

A uniform solid sphere has a radius of 1.50 m. An applied torque of 9.50 N.m gives the sphere an angular acceleration of 6.00 rad/s^2 about a fixed axis through its center. What is the mass of the sphere?

- A) 1.76 kg
- B) 1.06 kg
- C) 1.41 kg
- D) 2.11 kg
- E) 1.59 kg


Q13.

A particle is moving in a circle of radius 2.0 m with a tangential acceleration of 4.3 m/s^2 . At an instant when the magnitude of the total acceleration is 6.0 m/s^2 , what is the speed of the particle?

- A) 2.9 m/s
- B) 3.9 m/s
- C) 3.5 m/s
- D) 2.5 m/s
- E) 1.4 m/s

Q14.


Two equal particles, labeled **A** and **B** in **Figure 5**, are attached to a massless rod with a frictionless pivot at point **P**. The system is made to rotate clockwise with angular speed ω on a horizontal, frictionless tabletop. Particle **A** collides with and sticks to another equal particle that is at rest on the tabletop. What is the angular speed of the system immediately after the collision?



- A) 0.56ω
- B) 0.60ω
- C) ω
- D) 0.82ω
- E) 0.29ω

Q15.


A uniform beam has a weight of 120 N, and is supported as shown in **Figure 6**. What is the magnitude of the force by the pin on the beam?



- A) 75 N
- B) 94 N
- C) 88 N
- D) 63 N
- E) 90 N

Q16.

A traffic light hangs from the structure shown in **Figure 7**. The uniform rod AB is 4.50 m long and has a mass of 5.00 kg. The mass of the traffic light is 10.0 kg. Determine the magnitude of the tension in the horizontal massless cable CD.



- A) 121 N
- B) 160 N
- C) 91.0 N
- D) 100 N
- E) 145 N

Q17.

One end of a plastic rope, of length 45.0 m and radius 3.50 mm, is fixed to a ceiling while the other end is free. Its length increases by 1.10 m when a mass of 65.0 kg is attached to its free end. What is Young's modulus for plastic?

- A) $6.78 \times 10^8 \text{ N/m}^2$
- B) $4.69 \times 10^8 \text{ N/m}^2$
- C) $6.25 \times 10^8 \text{ N/m}^2$
- D) $2.83 \times 10^8 \text{ N/m}^2$
- E) $8.54 \times 10^8 \text{ N/m}^2$

Q18.

Two particles of mass M are initially separated by distance D . They are released from rest and accelerate toward each other through gravitational attraction. What is the kinetic energy of each particle when their separation is $D/3$?

- A) GM^2/D
- B) $3 GM^2/D$
- C) $GM/(2D^2)$
- D) $4 GM^2/D$
- E) $GM^2/2D$

Q19.

An object is released from rest at a height h above the surface of a planet of mass M and radius R . What is the speed of the object just before striking the surface of the planet? Take $h = 4000$ km, $R = 5000$ km and $M = 4.0 \times 10^{24}$ kg.

A) 6.9 km/s

B) 7.8 km/s


C) 3.5 km/s

D) 5.4 km/s

E) 4.8 km/s

Q20.

Three particles, each with a mass of 5.0 kg, are located at points in the xy plane as shown in **Figure 8**. What is the magnitude of the gravitational force on the particle at the origin due to the other two particles?



A) 2.1×10^{-8} N

B) 2.7×10^{-8} N


C) 1.8×10^{-8} N

D) 3.4×10^{-8} N

E) 2.9×10^{-8} N

Q21.

A planet moves around the Sun in the elliptical orbit shown in **Figure 9**. At point A, it is a distance of 1.75×10^8 km from the Sun and has a speed of 40 km/s. What is its speed at point B which is a distance of 2.50×10^8 km from the Sun?



A) 28 km/s

B) 11 km/s

C) 34 km/s

D) 40 km/s

E) 57 km/s


Q22.

The mass of a planet is $1/100$ that of Earth and its radius is $1/4$ that of Earth. If a person has a weight of 150 N on the surface of Earth, what would be his weight on the surface of the planet?

- A) 24 N
- B) 940 N
- C) 6.0 N
- D) 150 N
- E) 38 N

Q23.


Figure 10 shows a stream of water flowing through a hole at depth $h = 10\text{ cm}$ in a tank holding water to height $H = 50\text{ cm}$ and whose upper surface is open to the atmosphere. What is the speed of water exiting the hole?



- A) 1.4 m/s
- B) 2.3 m/s
- C) 3.5 m/s
- D) 1.8 m/s
- E) 2.9 m/s

Q24.

The density of oil is 0.80 g/cm^3 . What is the height h of the column of oil shown in **Figure 11**?



- A) 10 cm
- B) 12 cm
- C) 2.0 cm
- D) 4.6 cm
- E) 8.0 cm

Q25.

An object has a weight of 30 N in air. It has a weight of 25 N when completely submerged in water. What is the volume of the object?

- A) $5.1 \times 10^{-4} \text{ m}^3$
- B) $4.6 \times 10^{-4} \text{ m}^3$
- C) $3.1 \times 10^{-4} \text{ m}^3$
- D) $2.6 \times 10^{-4} \text{ m}^3$
- E) $2.0 \times 10^{-4} \text{ m}^3$


Q26.

Water, with a pressure of $3.5 \times 10^5 \text{ Pa}$, is flowing at a speed of 5.0 m/s in a horizontal pipe. The area of the pipe is reduced to 1/3 its original value. What are the pressure and the speed of the water after the reduction?

- A) $2.5 \times 10^5 \text{ Pa}$, 15 m/s
- B) $3.0 \times 10^5 \text{ Pa}$, 10 m/s
- C) $3.0 \times 10^5 \text{ Pa}$, 15 m/s
- D) $4.5 \times 10^5 \text{ Pa}$, 1.5 m/s
- E) $5.5 \times 10^5 \text{ Pa}$, 1.5 m/s

Q27.

A block of mass $M = 5.4 \text{ kg}$, at rest on a horizontal frictionless table, is attached to a rigid support by a spring of force constant $k = 6000 \text{ N/m}$ (see **Figure 12**). A bullet of mass $m = 9.5 \text{ g}$ and speed of 630 m/s strikes and is embedded in the block. What is the amplitude of the resulting harmonic motion?



- A) 3.3 cm
- B) 2.8 cm
- C) 1.3 cm
- D) 7.6 cm
- E) 3.8 cm

Q28.

A 0.50-kg mass connected to a spring is moving on a frictionless surface and its displacement is given by: $x(t) = 0.32 \cos(7.4t)$, where x is in meters and t is in seconds. What is the mechanical energy of the system?

- A) 1.4 J
- B) 0.90 J
- C) 0.47 J
- D) 4.4 J
- E) 0.19 J

Q29.

A block attached to a spring oscillates in simple harmonic motion along the x -axis with amplitude x_m . Its total energy is 50.0 J. What is its kinetic energy when $x = x_m/2$?

A) 37.5 J

B) 12.5 J

C) 25.0 J

D) 50.0 J

E) zero

Q30.

A simple pendulum has a frequency of 3 Hz. To increase its frequency to 6 Hz

A) Decrease its length by a factor of 4

B) Decrease its length by a factor of 2

C) Increase its length by a factor of 4

D) Increase its length by a factor of 2

E) Decrease its mass by a factor of 4
