

Q1.

The velocity of a particle is given by  $v = At^2 + (B/A)t$ , where  $v$  is in m/s and  $t$  is in seconds. The dimension of  $B$  is:

- A)  $L^2 T^{-5}$
- B)  $L^3 T^{-3}$
- C)  $L^2 T^3$
- D)  $L^4 T^4$
- E)  $L^5 T^{-6}$

Q2.


The position of a particle as a function of time is given by  $x(t) = 3.0t^3 - 10.0t^2 + 9.0t$ , where  $x$  is in meters and  $t$  is in seconds. Find the velocity at the time when the acceleration is zero.

- A)  $-2.1$  m/s
- B)  $+2.1$  m/s
- C)  $+5.0$  m/s
- D)  $-5.0$  m/s
- E)  $+3.5$  m/s

Q3.

The position-time curve for a particle moving along the  $x$ -axis is shown in **Figure 1**. The dashed straight lines are tangent to the curve at  $t = 1.0$  s and  $t = 3.0$  s. Find the magnitude of the average acceleration in the time interval  $t = 1.0$  s and  $t = 3.0$  s.

Fig#



- A)  $1.5$  m/s<sup>2</sup>
- B)  $5.0$  m/s<sup>2</sup>
- C)  $1.0$  m/s<sup>2</sup>
- D)  $2.0$  m/s<sup>2</sup>
- E)  $2.5$  m/s<sup>2</sup>

Q4.

Two objects A and B are thrown from the top of a building with the same magnitude of velocity. Object A is thrown upward but object B is thrown downward. When they reach the ground (ignore air resistance):

- A) The two objects have the same velocity.
- B) Object A has a higher velocity than object B.
- C) Object B has a higher velocity than object A.
- D) The velocities will depend on the masses of the objects.
- E) The velocities will depend on the shapes of the objects.

---

Q5.

A ball starts from rest and slides down a hill with a constant acceleration for 10.0 s. If it travels 50.0 m during the first 5.00 seconds of its motion, how far will it travel during the next 5.00 seconds of its motion?


- A) 150 m
- B) 98.0 m
- C) 75.0 m
- D) 200 m
- E) 25.0 m

---

Q6.

**Figure 2** shows three vectors **A**, **B** and **C**, whose magnitudes are 15 m, 10 m and 12 m respectively. Find both the magnitude and direction of the resultant of these vectors.

Fig#



- A) 18.0 m,  $129^\circ$  from the positive x-axis
  - B) 14.0 m,  $150^\circ$  from the positive x-axis
  - C) 12.0 m,  $53^\circ$  from the positive x-axis
  - D) 20.0 m,  $163^\circ$  from the positive x-axis
  - E) 10.0 m,  $110^\circ$  from the positive x-axis
-

Q7.

If vector  $\mathbf{A} = b(3.0\mathbf{i} + 4.0\mathbf{j})$ , where  $b$  is a constant. Find the value of  $b$  that makes vector  $\mathbf{A}$  a unit vector.

- A) 0.20
- B) 0.10
- C) 0.30
- D) 0.50
- E) 0.40

---

Q8.

Two vectors  $\vec{A}$  and  $\vec{B}$  have magnitudes 3.0 and 4.0 respectively. Their vector product is  $\vec{A} \times \vec{B} = -5.0\vec{k} + 2.0\vec{i}$ . Find the angle between  $\vec{A}$  and  $\vec{B}$ .

- A)  $27^\circ$
- B)  $22^\circ$
- C)  $39^\circ$
- D)  $11^\circ$
- E)  $16^\circ$

---

Q9.


A player running in an open field has components of velocity  $v_x = 2.60$  m/s and  $v_y = -1.80$  m/s at  $t = 10.0$  s. For the time interval from  $t = 10.0$  s to  $t = 20.0$  s, the magnitude of its average acceleration is  $0.45$  m/s<sup>2</sup> and makes an angle of  $31.0^\circ$  with the positive x-axis. Find the velocity (in units m/s) of the player at  $t = 20.0$  s.

- A)  $6.46\mathbf{i} + 0.518\mathbf{j}$
  - B)  $5.32\mathbf{i} + 3.52\mathbf{j}$
  - C)  $4.52\mathbf{i} + 5.60\mathbf{j}$
  - D)  $2.46\mathbf{i} + 2.52\mathbf{j}$
  - E)  $9.60\mathbf{i} + 5.30\mathbf{j}$
-

Q10.

A basketball player would like to throw a ball at an angle of  $\theta_0 = 60^\circ$  above the horizontal such that the ball just goes through the center of the rim of the basket that is  $h_2 = 3.0$  m high from the floor and it is at a horizontal distance of  $d_1 = 5.0$  m from the player's hand (see **Figure 3**). At the instant the ball leaves the player's hand, his hand is  $h_1 = 2.0$  m above the floor. Find the magnitude of the initial velocity of the ball.

Fig#



- A) 8.0 m/s
- B) 5.0 m/s
- C) 9.8 m/s
- D) 3.2 m/s
- E) 7.0 m/s

Q11.

A particle rotates clockwise with a constant speed in a horizontal circle whose center is at the origin. It completes one revolution in 2.0 s. At  $t = 0.0$ , the particle is at  $(0.0, 1.5)$  m. What is the magnitude of the average acceleration of the particle in the interval between  $t = 3.0$  and  $t = 8.0$  seconds?

- A)  $1.9 \text{ m/s}^2$
- B)  $4.7 \text{ m/s}^2$
- C) 0
- D)  $1.0 \text{ m/s}^2$
- E)  $0.60 \text{ m/s}^2$

Q12.

A 100-m wide river flows due east at a uniform speed of 3.0 m/s. A boat with a speed of 8.0 m/s relative to the water leaves the south bank pointed in a direction  $60^\circ$  west of north. What is the boat's velocity relative to the ground?

[ $\mathbf{i}$  and  $\mathbf{j}$  are unit vectors along the east and north directions respectively]

- A)  $-3.9 \mathbf{i} + 4.0 \mathbf{j}$  m/s
- B)  $-3.0 \mathbf{i} + 8.0 \mathbf{j}$  m/s
- C)  $-3.0 \mathbf{i} + 4.0 \mathbf{j}$  m/s
- D)  $+11 \mathbf{i} + 5.0 \mathbf{j}$  m/s
- E)  $-11 \mathbf{i} + 5.0 \mathbf{j}$  m/s

---

Q13.

Which one of the following statements is CORRECT?


- A) It is possible to be accelerating while traveling at constant speed.
- B) It is not possible to be accelerating while traveling at constant speed.
- C) It is possible to round a curve with zero acceleration.
- D) It is not possible to round a curve with a constant magnitude of acceleration.
- E) The acceleration of a particle moving in a circle with constant speed is constant.

---

Q14.

As shown in **Figure 4**, a man is pulling a 60.0 N box up an inclined frictionless plane with a force  $F = 23.7$  N whose direction makes an angle  $30.0^\circ$  with inclined plane. The inclined plane makes an angle of  $20.0^\circ$  with the horizontal. Find the magnitude of the acceleration of the box.

Fig#




- A) Zero
  - B)  $3.35 \text{ m/s}^2$
  - C)  $9.80 \text{ m/s}^2$
  - D)  $4.80 \text{ m/s}^2$
  - E)  $6.50 \text{ m/s}^2$
-

Q15.

Two boxes A and B, are connected to the ends of a light vertical cord, as shown in **Figure 5**. A constant upward force  $F = 80.0 \text{ N}$  is applied to box A. Starting from rest, box B descends  $12.0 \text{ m}$  in  $4.00 \text{ s}$ . Find the mass of box B if the tension in the cord is  $36.0 \text{ N}$ .

Fig#



- A) 4.34 kg
- B) 3.42 kg
- C) 24.0 kg
- D) 15.3 kg
- E) 12.7 kg

---

Q16.

With what force will the feet of a person of mass  $60.0 \text{ kg}$  press downward on an elevator floor when the elevator has an upward acceleration of  $1.20 \text{ m/s}^2$ ?

- A) 660 N
- B) 600 N
- C) 516 N
- D) 588 N
- E) 980 N

---

Q17.


A  $0.15 \text{ kg}$  ball is thrown at an angle of  $30^\circ$  above the horizontal with an initial speed of  $12 \text{ m/s}$ . At its highest point, the net force on the ball is:

- A) 1.5 N, down
  - B) 9.8 N,  $30^\circ$  below horizontal
  - C) 0
  - D) 9.8N, up
  - E) 9.8 N, down
-

Q18.

As shown in **Figure 6**, blocks  $m_1$  and  $m_2$  have masses of 4.00 kg and 8.00 kg, respectively. The coefficient of kinetic friction between  $m_1$  and the horizontal surface is 0.500. The incline plane ( $\theta = 30^\circ$ ) is frictionless. Find the acceleration of the system. (Assume that the pulley is massless and frictionless).

Fig#




- A)  $1.63 \text{ m/s}^2$
- B)  $3.48 \text{ m/s}^2$
- C)  $4.80 \text{ m/s}^2$
- D) 0
- E)  $0.430 \text{ m/s}^2$

Q19.

**Figure 7** shows an object of mass  $m = 0.10 \text{ kg}$  tied to a rope rotating in a horizontal circle of radius  $r = 0.25 \text{ m}$ , on a frictionless table top. It rotates at constant speed of  $4.0 \text{ m/s}$  while the mass  $M$  is stationary. Find the value of mass  $M$ .

Fig#




- A) 0.65 kg
- B) 0.87 kg
- C) 0.25 kg
- D) 0.98 kg
- E) 1.0 kg

Q20.

**Figure 8** shows a constant horizontal force  $F = 20.0 \text{ N}$  applied to block A (mass =  $2.00 \text{ kg}$ ) which pushes against block B (mass =  $3.00 \text{ kg}$ ) to the right. The coefficient of kinetic friction between the surface and object A is  $0.220$  and between the surface and object B is  $0.350$ . Find the magnitude of the contact force between object A and object B.

Fig#



- A)  $13.5 \text{ N}$
  - B)  $20.0 \text{ N}$
  - C)  $5.40 \text{ N}$
  - D)  $11.4 \text{ N}$
  - E)  $15.0 \text{ N}$
-