

Q1.

A hectare is a unit of area that is equal to $1.0 \times 10^4 \text{ m}^2$. If water of volume 0.020 km^3 covers 30 hectares of flat land, find the depth of the water.

- A) 67 m
- B) 26 m
- C) 45 m
- D) 30 m
- E) 87 m

Q2.


Consider the following equation: $x = At^2 + \frac{B}{(v + \alpha)}t$, where x is the distance, t is the time and v is the speed. Find the dimensions of B :

- A) L^2T^{-2}
- B) L^2T
- C) LT^{-1}
- D) LT^2
- E) L

Q3.

Figure 1 gives the acceleration of a particle as a function of time. In which of the time intervals indicated does the particle move with constant speed?

- A) E
- B) C, G
- C) C, E, G
- D) A, B, H
- E) D, F,



Q4.

At time $t = 0$, a particle had a speed of 20 m/s in the positive x direction. At time $t = 2.5 \text{ s}$, its speed was 40 m/s in the opposite direction. Find the average acceleration of the particle during the 2.5 s interval.

- A) -24 m/s^2
- B) $+18 \text{ m/s}^2$
- C) -8.0 m/s^2
- D) $+20 \text{ m/s}^2$
- E) -30 m/s^2

Q5.

A car travels in a straight line. First, it starts from rest at point A and accelerates at a rate of 5.00 m/s^2 until it reaches a speed of 100 m/s at point B. The car then slows down at a constant rate of 8.00 m/s^2 until it stops at point C. Find the time the car takes for this trip (from point A to point C).

- A) 32.5 s**
- B) 25.0 s
- C) 10.5 s
- D) 15.0 s
- E) 45.0 s

Q6.

A parachutist jumps from an airplane at an altitude of $5.0 \times 10^3 \text{ m}$. He falls with an acceleration $g = 9.8 \text{ m/s}^2$ for the first 10 s . Then he opens his parachute and falls with a net vertical upward acceleration of 50 m/s^2 until his downward speed reaches 20 m/s . How far does he fall vertically downward when his net upward acceleration was 50 m/s^2 ?

- A) 92 m**
- B) 50 m
- C) 75 m
- D) 67 m
- E) 45 m

Q7.


Two vectors are given by $\vec{A} = 2.00\hat{i} + 2.00\hat{j}$ and $\vec{B} = -2.00\hat{i} + 4.00\hat{j}$, find the angle between \vec{A} and \vec{B} .

- A) 71.6°**
 - B) 45.0°
 - C) 56.1°
 - D) 18.4°
 - E) 24.5°
-

Q8.

The two vectors shown in **Figure 2** lie in an xy plane. What are the signs of the x and y components, respectively, of the vector $(\vec{d}_2 - \vec{d}_1)$?

- A) +, +
- B) +, -
- C) -, +
- D) -, -
- E) None of the other answers is correct.



Q9.

For the following three vectors, find $\vec{C} \cdot (2\vec{A} \times \vec{B})$

$$\vec{A} = 2.00\hat{i} + 3.00\hat{j}$$

$$\vec{B} = -3.00\hat{i} + 4.00\hat{j}$$

$$\vec{C} = 7.00\hat{i} + 3.00\hat{k}$$

- A) 102
- B) -14.0
- C) 0
- D) 56.0
- E) 78.0

Q10.

A man makes three successive displacements; 3.50 m south, 8.20 m northeast, and 15.0 m west, respectively. Find the resultant displacement (both the magnitude and direction relative to the east and measured counter-clock wise).

- A) 9.48 m, 166°
- B) 9.48 m, 45.0°
- C) 9.48 m, 225°
- D) 5.80 m, 45.0°
- E) 5.80 m, 225°

Q11.

A ship sails due north at 4.50 m/s relative to the ground while a boat heads northwest with a speed of 5.20 m/s relative to the ground. Find the speed of the ship relative to the boat.

- A) 3.77 m/s**
- B) 2.39 m/s
- C) 7.95 m/s
- D) 1.25 m/s
- E) 6.11 m/s

Q12.

A student throws a red ball from the balcony of a tall building with an initial horizontal speed of 10 m/s. At the same time, a second student drops a blue ball from the same balcony. Neglecting air resistance, which statement is true?

- A) The two balls reach the ground at the same instant.**
- B) The blue ball reaches the ground first.
- C) The red ball reaches the ground first.
- D) Both balls hit the ground with the same speed.
- E) The blue ball hits the ground with larger speed.

Q13.

A stone is tied to the end of a string and is rotated in a horizontal circle at 400 revolutions per minute. If the magnitude of its acceleration is $1.5 \times 10^3 \text{ m/s}^2$, what is the radius of the circle?

- A) 0.85 m**
- B) 0.35 m
- C) 0.64 m
- D) 0.71 m
- E) 0.53 m

Q14.

A ball is thrown straight upward and returns to the thrower's hand (at the same initial level) after 3.00 s. A second ball thrown from the same height at an angle of 37.0° with the horizontal reaches the same maximum height as the first ball. With what speed was the second ball thrown?

- A) 24.4 m/s**
 - B) 14.7 m/s
 - C) 29.1 m/s
 - D) 49.3 m/s
 - E) 35.2 m/s
-

Q15.

A particle starts from the origin of an xy plane. Its acceleration is given by $\vec{a} = (2.0\hat{i} + 4.0\hat{j}) \text{ m/s}^2$. At time $t = 0$, the velocity is $-4.0\hat{i} \text{ m/s}$. What is the particle's velocity if the y-component of its displacement is $+18 \text{ m}$?

- A) $(2.0\hat{i} + 12\hat{j}) \text{ m/s}$
 - B) $(4.0\hat{i} - 6.0\hat{j}) \text{ m/s}$
 - C) $(2.0\hat{i} + 2.0\hat{j}) \text{ m/s}$
 - D) $(3.0\hat{i} + 12\hat{j}) \text{ m/s}$
 - E) $(4.0\hat{i} - 4.0\hat{j}) \text{ m/s}$
-