

Q1.

The air resistance force on a falling object can be expressed as $F = a v^2$, where a is a constant, and v is the speed of the object. The dimension of a is

- A) M/L
- B) ML
- C) L/M
- D) ML^2
- E) ML^2


Q2.

Assume it takes 6.00 minutes to fill a 30.0-gallon tank. Calculate the rate at which the tank is filled in cubic meters per second. [1 gallon = 231 inch³, 1 inch = 2.54 cm]

- A) 3.15×10^{-4}
- B) 4.89×10^{-5}
- C) 5.25×10^{-5}
- D) 1.89×10^{-2}
- E) 1.05×10^{-5}

Q3.

The top diagram in **Figure 1** represents a series of the locations of a particle moving along a straight line from left to right. The dots are taken every one second. Which of the lower graphs represents the motion of the particle?



- A) IV
- B) I
- C) II
- D) III
- E) V

Q4.

The position of a particle moving along the x axis is given by: $x(t) = 1.5 t^2 - 0.050 t^3$, where x in meters and t is in seconds. Calculate the average acceleration of the particle during the interval from $t = 2.0$ s to $t = 4.0$ s.

- A) 2.1 m/s²
- B) 1.7 m/s²
- C) 0.45 m/s²
- D) 9.6 m/s²
- E) 5.4 m/s²


Q5.

A car travels in a straight line a distance of 40 m in 8.0 s while slowing down at constant deceleration to a final speed of 2.5 m/s. Find its initial speed.

- A) 7.5 m/s
- B) 13 m/s
- C) 2.5 m/s
- D) 4.2 m/s
- E) 6.8 m/s

Q6.

A rock is thrown vertically upward from point A at the roof of a building (see **Figure 2**). It reaches point B, which is 30.0 m below point A, in a time of 5.00 s after it is thrown. What is the initial speed of the rock? Ignore air resistance.



- A) 18.5 m/s
- B) 30.5 m/s
- C) 24.2 m/s
- D) 49.0 m/s
- E) 39.8 m/s

Q7.

If two vectors have the same magnitude, what should be the angle between them for their resultant to have the same magnitude as any of them?

- A) 120°
- B) 60°
- C) 45°
- D) 30°
- E) 150°

Q8.

A person moves 180 m straight west, then 270 m at 30.0° east of north. What third displacement would bring him back to the starting point?

- A) 238 m at 79.1° south of east
- B) 392 m at 10.9° north of east
- C) 194 m at 25.7° west of north
- D) 169 m at 29.3° west of south
- E) 248 m at 36.3° east of south

Q9.

Two vectors \vec{A} and \vec{B} lie in the xy planes. Their magnitudes and angles measured counterclockwise from the positive x -axis are: $A = 5.0$, $\theta_A = 58^\circ$, $B = 4.0$, $\theta_B = 28^\circ$. A third vector \vec{C} has magnitude 6.0 and points along the positive z -axis. Find $(\vec{B} \times \vec{A}) \cdot \vec{C}$.

- A) + 60
- B) - 34
- C) - 60
- D) zero
- E) + 34


Q10.

The position vector (in meters) of a particle is given by $\vec{r} = 2.50 t^2 \hat{i} + 5.00 t \hat{j}$, where t is in seconds. At $t = 2.00$ s, what is the instantaneous speed (v) of the particle and the angle θ between \vec{v} and the positive x axis measured counterclockwise?

- A) $v = 11.2$ m/s, $\theta = 26.6^\circ$
- B) $v = 11.2$ m/s, $\theta = 63.4^\circ$
- C) $v = 14.1$ m/s, $\theta = 26.6^\circ$
- D) $v = 14.1$ m/s, $\theta = 63.4^\circ$
- E) $v = 12.6$ m/s, $\theta = 45.0^\circ$

Q11.


A small stone is thrown with an initial speed of 6.5 m/s at an angle of 60° above the horizontal and lands on a shelf that is a horizontal distance of 2.5 m from its launch point (see **Figure 3**). What is the height (h) of the shelf? Ignore air resistance.



- A) 1.4 m
- B) 4.3 m
- C) 5.7 m
- D) 3.6 m
- E) 2.9 m

Q12.

A particle executes uniform circular motion with it moves clockwise with a speed of 5.00 m/s around a circle of radius 50.0 m, as shown in **Figure 4**. What is the least time to go from point A to point B?



- A) 15.7 s
- B) 62.8 s
- C) 31.4 s
- D) 47.1 s
- E) 39.2 s

Q13.

A car has a velocity of 15 m/s due south as it passes a train travelling with a velocity of 24 m/s due north. What is the velocity of the car relative to the train?

- A) 39 m/s, due south
- B) 39 m/s, due north
- C) 9 m/s, due south
- D) 9 m/s, due north
- E) 15 m/s, due north


Q14.

Two cars A and B approach each other at an intersection. Car A is travelling due south at 20 m/s, while car B is travelling due east at 17 m/s. What is the speed of car A relative to car B?

- A) 26 m/s
- B) 37 m/s
- C) 11 m/s
- D) 21 m/s
- E) 24 m/s

Q15.

A box of weight W hangs from two massless strings, as shown in **Figure 5**. Each string makes the same angle θ with the horizontal. The magnitudes of the weight of the box and tension in each string are equal ($T = W$) if the angle θ is



- A) 30°
- B) 15°
- C) 45°
- D) 60°
- E) 75°

Q16.

A 4.8-kg box is pulled vertically upward with a tension of 72 N. What is the magnitude of the acceleration of the box?

- A) 5.2 m/s^2
- B) 25 m/s^2
- C) 1.1 m/s^2
- D) 2.7 m/s^2
- E) 6.7 m/s^2


Q17.

A 2.50-kg object is subject to the gravitational force and another constant force. The object starts from rest and in 2.00 s experiences a displacement of $(3.00\hat{i} - 3.50\hat{j})$ (m), where the direction of \hat{j} is the upward vertical direction. Determine the other force.

- A) $3.75\hat{i} + 20.1\hat{j}$ (N)
- B) $3.75\hat{i} - 4.38\hat{j}$ (N)
- C) $3.75\hat{i} + 32.3\hat{j}$ (N)
- D) $3.75\hat{i} - 32.3\hat{j}$ (N)
- E) $3.75\hat{i} - 24.5\hat{j}$ (N)

Q18.


Three blocks are in contact with one another on a frictionless horizontal surface, as shown in **Figure 6**. Take $m_1 = 3.00$ kg, $m_2 = 4.00$ kg, and $m_3 = 5.00$ kg. A horizontal force \vec{F} , of magnitude 18.0 N, is applied to m_1 as shown. What is the magnitude of the contact force between blocks m_1 and m_2 ?



- A) 13.5 N
- B) 4.50 N
- C) 22.5 N
- D) 6.00 N
- E) 11.6 N

Q19.

As shown in **Figure 7**, block A (mass 2.3 kg) rests on a horizontal rough surface ($\mu_k = 0.45$). It is connected by a horizontal cord passing over a massless frictionless pulley to block B (mass 1.3 kg). What is the magnitude of the acceleration of the system?



- A) 0.72 m/s^2
- B) 0.15 m/s^2
- C) 0.65 m/s^2
- D) 0.38 m/s^2
- E) 0.34 m/s^2

Q20.

A small car of mass 0.750 kg travels at constant speed on the inside of a track that is a vertical circle, as shown in **Figure 8**. If the normal force exerted by the track on the car when it is at the top of the track (point B) is 5.50 N , what is the magnitude of the normal force at the bottom of the track (point A)?



- A) 20.2 N
 - B) 9.20 N
 - C) 7.40 N
 - D) 14.7 N
 - E) 12.9 N
-
