


Q1. Two persons pushed a car initially at rest at its front doors, each applying a force with magnitude $|\vec{F}| = 300 \text{ N}$ at 25.0° to the forward direction, as shown in **Figure 1**. How much average power does **each person** requires in pushing the car 10.0 m for 10.0 seconds?
Fig#




- A) 272 W
- B) 145 W
- C) 710 W
- D) 424 W
- E) 299 W

Q2. A spring has a spring constant k . If the work done in stretching the spring a distance $x = L$ from the equilibrium position is W , the work required to stretch the spring from $x_i = L$ to $x_f = 2L$ will be:

- A) 3 W
- B) 5 W
- C) 4 W
- D) 2 W
- E) 1 W


Q3. A single force acts on the body causing the body to move in a straight line. A plot of the body's velocity v (m/s) versus time t (s) is shown in the Fig 2. The **correct** statement among the following is:
Fig#



- A) in moving from D to E, the work done by the force on the body is positive.
- B) in moving from B to C no work is done on the body but the body does work on the system.
- C) in moving from C to D, the work done by the force on the body is positive.
- D) in moving from A to B, the work done by the force on the body is negative.
- E) in moving from A to D, the work done by the force on the body is positive.


Q4. A block, of mass $m = 200 \text{ g}$, slides back and forth on a **frictionless surface** between two springs, as shown in **Figure 3**. The left-hand side spring has $k_1 = 130 \text{ N/m}$ and its maximum compression is 16 cm . The right-hand side spring has $k_2 = 280 \text{ N/m}$. Find the maximum compression of the right-hand side spring.

Fig#



- A) 11 cm.
- B) 14 cm.
- C) 2.0 cm.
- D) 30 cm.
- E) 8.0 cm.

Q5. A block, of mass $m = 1.0 \text{ kg}$, initially at rest, falls from a height of $h = 0.50 \text{ m}$, on a vertical spring fastened to a horizontal board placed on the floor, as shown in **Figure 4**. If the spring constant is $k = 98 \text{ N/m}$, the maximum compression that the spring undergoes is:
Fig#




- A) 0.43 m
- B) 0.17 m
- C) 0.34 m
- D) 0.86 m
- E) 0.54 m

Q6. A single force F , of magnitude 10.0 N , accelerates an object of mass 5.00 kg for three seconds starting from rest at $t = 0$. What is the work done on the object in the time interval from $t = 2.00 \text{ sec}$ to $t = 3.00 \text{ sec}$.

- A) 50.0 J.
- B) 40.0 J.
- C) 20.0 J.
- D) 10.0 J.
- E) 25.0 J.


Q7. A block slides back and forth in a hemispherical bowl, starting from rest at the top point **A**, as shown in **Figure 5**. The bowl is frictionless except for a 2.0 cm -wide rough flat surface at the bottom, where coefficient of kinetic friction is $\mu_k = 0.45$. How many times does the block cross the rough region before coming to rest?
Fig#



- A) 10 times.
- B) 14 times.
- C) 13 times.
- D) 3 times.
- E) 4 times.

Q8. **Figure 6** shows a block, of mass $m = 0.50 \text{ kg}$ with an initial speed of $v_i = 10 \text{ m/s}$, moving down an inclined rough plane of angle 30° . The coefficient of kinetic friction between the block and the plane is $\mu = 0.20$. The speed, v_f , of the block after it travels a distance of 10 m is:


Fig#



- A) 13 m/s
- B) 24 m/s
- C) 8 m/s
- D) 17 m/s
- E) 36 m/s

Q9. A force $F(t)$ (pointing in the + x-direction) acts on a ball with mass $m = 0.060$ kg initially at rest at $t = 1.00$ ms. **Figure 7** shows a plot of $F(t)$ vs t . Find the speed of the ball at $t = 2.50$ ms.

Fig#




- A) 225 m/s
- B) 375 m/s
- C) 400 m/s
- D) 153 m/s
- E) 642 m/s

Q10. A system consists of two particles m_1 and m_2 , where the mass of $m_2 = 0.10$ kg. At $t = 0$ s, the particle m_1 was at $x_1 = 0.0$ m and has a velocity \vec{v}_1 , and the other particle m_2 was at rest at $x_2 = 8.0$ m. At $t = 0$ s, the center of mass of the system was at $x_{com} = 2.0$ m, and has a velocity of $\vec{v}_{com} = 5.0 \hat{i}$ m/s. What was the velocity \vec{v}_1 ?

- A) $\vec{v}_1 = 6.7 \hat{i}$ m/s
- B) $\vec{v}_1 = 2.7 \hat{i}$ m/s
- C) $\vec{v}_1 = 1.8 \hat{i}$ m/s
- D) $\vec{v}_1 = 3.4 \hat{j}$ m/s
- E) $\vec{v}_1 = 9.2 \hat{i}$ m/s

Q11. After a completely inelastic collision between two objects of equal mass m , each having an initial speed v , the two move off together with a speed $v/3$, see **Figure 8**. What was the angle (2θ) between their initial velocities?

Fig#



- A) 141°
- B) 134°
- C) 102°
- D) 163°
- E) 127°

Q12. A ball of mass m_1 makes a head on elastic collision with second ball, of mass m_2 , initially at rest. If m_1 rebounds in the opposite direction with a speed equal to one-fourth its original speed, what is the mass m_2 ?

- A) $\frac{5}{3}m_1$
- B) $\frac{1}{2}m_1$
- C) $\frac{1}{3}m_1$
- D) $\frac{3}{4}m_1$
- E) $\frac{7}{2}m_1$

Q13. A uniform disk of 1.0 m radius is rotating about its symmetry axis with a constant angular speed of 2.0 rad/s. What are the magnitude of the tangential acceleration a_t and centripetal acceleration a_r of a point on the rim of the disk?

- A) $a_t = 0.0 \text{ m/s}^2$, $a_r = 4.0 \text{ m/s}^2$
- B) $a_t = 1.0 \text{ m/s}^2$, $a_r = 2.0 \text{ m/s}^2$
- C) $a_t = 0.0 \text{ m/s}^2$, $a_r = 1.0 \text{ m/s}^2$
- D) $a_t = 2.0 \text{ m/s}^2$, $a_r = 4.0 \text{ m/s}^2$


E) $a_t = 4.0 \text{ m/s}^2$, $a_r = 0.0 \text{ m/s}^2$

Q14. A disk, subjected to a constant net torque, rotates around a fixed axis starting from rest. The ratio of work done by the torque during the (0 – 5.0 s) interval to the work done during the (5.0 s-10 s) interval is:

- A) 1/3
- B) 1/2
- C) 3
- D) 2
- E) 4

Q15. A uniform disk, a thin hoop, and a uniform solid sphere, all with the same mass and same outer radius, are each free to rotate about a fixed axis through their centers. Identical forces are simultaneously applied to the rims of the objects, as shown in **Figure 9**. If the objects start from rest, rank the objects according to their angular speeds achieved after a given time (t sec), **least to greatest**.

Fig#




- A) hoop, disk, sphere
- B) All tie.
- C) hoop, sphere, disk
- D) disk, hoop, sphere
- E) sphere, disk, hoop

Q16. An engine applies a constant torque of 5.00 N·m on a wheel, with moment of inertia $I_0 = 10.0 \text{ kg}\cdot\text{m}^2$, to rotate it about its symmetry axis O. How much power is required by the engine to rotate the wheel at $t = 5.00 \text{ s}$, if the wheel starts from rest?

- A) 12.5 W
- B) 3.06 W
- C) 6.20 W
- D) 2.53 W
- E) 1.62 W

Q17. A single force F acts on a particle P . Rank each of the orientations of the force shown in **Figure 10** according to the magnitude of the time rate of change of the particle's angular momentum about the point O , **least to greatest**.


Fig#



- A) 1 and 2 tie, then 4, 3
- B) 1, 2, 3, 4
- C) 1 and 2 tie, then 3, 4
- D) 1 and 2 tie, then 3 and 4 tie
- E) All are the same

Q18. A disk with radius of 3.0 m and a moment of inertia of $8000\text{ kg}\cdot\text{m}^2$ is rotating about its central axis without friction with an angular velocity of 1.2 rad/s . Initially a man with mass m is moving with a velocity of 5.0 m/s , on a line tangent to the edge of the disk, as shown in **Figure 11**. The man jumps onto the edge of the disk. The final angular velocity of the disk and the man is 1.24 rad/s . The mass of the man m is:

Fig#



- A) 83 kg
- B) 61 kg
- C) 75 kg
- D) 53 kg
- E) 94 kg

Q19. A disk starts from rest and rotates with a constant angular acceleration. If the angular velocity is ω rad/s at the end of the first two revolutions, then at the end of the first eight revolutions, the angular velocity will be:

- A) 2ω rad/s
- B) $\sqrt{2} \omega$ rad/s
- C) 3ω rad/s
- D) 4ω rad/s
- E) 5ω rad/s

Q20. A hoop rolls without sliding along a horizontal floor. The ratio of its translational kinetic energy to its rotational kinetic energy (about an axis through its center of mass) is:

- A) 1
- B) 2
- C) 3
- D) $1/2$
- E) $1/3$