

Q1.

A 3.00×10^3 N automobile accelerates from rest to 50.0 m/s in 6.00 s with a constant acceleration. Calculate the instantaneous power delivered by the engine at $t = 6.00$ s.

- A) 12.8×10^4 W
- B) 15.0×10^4 W
- C) 9.45×10^5 W
- D) 24.5×10^3 W
- E) 4.51×10^7 W

Q2.

A man pushes a 30.0 kg box a horizontal distance of 4.50 m along a level floor at a constant velocity. The coefficient of kinetic friction between the box and the floor is 0.250. Find the total work done on the box.

- A) 0
- B) 74.0 J
- C) 333 J
- D) -330 J
- E) 350 J

Q3.


A 5.0 kg box is taken from point A (1.0, 2.0, -2.0) m to point B (6.0, -3.0, -2.0) m by applying a single constant force $\mathbf{F} = (3.0 \text{ N}) \mathbf{i} + (2.0 \text{ N}) \mathbf{j} + (5.0 \text{ N}) \mathbf{k}$. Find the change in the kinetic energy of the box.

- A) 5.0 J
 - B) 25 J
 - C) 10 J
 - D) 18 J
 - E) 2.0 J
-

Q4.

Figure 1 shows a plot of the acceleration a_x versus the displacement x for a particle of mass $m = 2.0$ kg moving along the x -axis. The scale of the figure's vertical axis is set by $a_s = 3.0$ m/s^2 . How much work is done on the particle as it moves from $x = 2.0$ to $x = 6.0$ m.

Fig#




- A) 12 J
- B) 9.0 J
- C) 6.0 J
- D) 18 J
- E) 24 J

Q5.

A massless spring has a spring constant of 500 N/m. A 2.0 kg object is released from rest at a height $h = 1.0$ m above the spring and lands on it (**Figure 2**). Find the object's speed when the spring is compressed 20 cm.

Fig#




- A) 3.7 m/s
- B) 3.1 m/s
- C) 4.1 m/s
- D) 4.5 m/s
- E) 4.9 m/s

Q6.

A block with mass $m = 0.50$ kg is forced against a horizontal spring of spring constant 100 N/m and negligible mass, compressing the spring a distance of 0.20 m (**Figure 3**). When released, the block moves on a horizontal tabletop for 1.0 m before coming to rest. Find the coefficient of kinetic friction μ_k between the block and the tabletop.

Fig#



- A) 0.41
- B) 0.35
- C) 0.25
- D) 0.13
- E) 0.53

Q7.


The work done by a conservative force acting on a body (Choose the CORRECT answer):

- A) Does not change the total energy.
 - B) Does not change the potential energy.
 - C) Is always equal to zero.
 - D) Does not change the kinetic energy.
 - E) Is always equal to the sum of the changes in potential and kinetic energy.
-

Q8.

A 15.0 kg stone slides down a smooth snow-covered hill (**Figure 4**), leaving point A with a speed of 4.0 m/s. Then it slides a distance of 118 m on a rough horizontal surface from point B to point C before coming to rest. Find the coefficient of kinetic friction μ_k between the stone and the surface.

Fig#



- A) 0.600
- B) 0.500
- C) 0.400
- D) 0.550
- E) 0.450

Q9.

Two particles of masses 3.0 kg and 5.0 kg are moving with initial velocities of $(-3.0 \mathbf{i} + 4.0 \mathbf{j})$ m/s and $(2.0 \mathbf{i} + 3.0 \mathbf{j})$ m/s respectively. They collide completely inelastically. Find the velocity of the center of mass of the two particles after the collision.

- A) $(0.13 \mathbf{i} + 3.4 \mathbf{j})$ m/s
- B) $(3.2 \mathbf{i} + 4.4 \mathbf{j})$ m/s
- C) $(5.13 \mathbf{i} + 1.34 \mathbf{j})$ m/s
- D) $(-9.00 \mathbf{i} + 12.0 \mathbf{j})$ m/s
- E) $(10.0 \mathbf{i} + 15.0 \mathbf{j})$ m/s

Q10.


A car with a mass of 1.2×10^3 kg is travelling to the right at a speed of 15 m/s when it collides head-on with a truck of mass 2.0×10^3 kg travelling at a speed of 15 m/s to the left. The vehicles lock together when they collide. Find the average force (both magnitude and direction) exerted on the car if the collision lasts for 0.20 s.

- A) 1.1×10^5 N to the left
- B) 1.1×10^5 N to the right
- C) 2.2×10^4 N to the left
- D) 3.1×10^4 N to the right
- E) 5.3×10^5 N to the left

Q11.

Two objects A and B, with the same mass collide on ice with negligible friction. **Figure 5** gives speeds and directions of the objects BEFORE and AFTER the collision. Find the speed v and angle θ for object A after the collision.

Fig#




- A) 5.0 m/s , 37°
 - B) 7.0 m/s , 45°
 - C) 10 m/s, 30°
 - D) 3.5 m/s, 50°
 - E) 1.4 m/s, 20°
-

Q12.

A machine part consists of three objects welded together: A) a thin, uniform 4.00 kg bar that is 1.50 m long, B) a vertical bar of mass 3.00 kg and length 1.80 m and C) dense 2.00 kg ball attached to the end of object B (**Figure 6**). Find the center of mass of this system.

Fig#




- A) $(-0.333\text{m}, -0.700\text{ m})$
- B) $(-0.750\text{m}, -0.900\text{ m})$
- C) $(-0.250\text{m}, -0.450\text{ m})$
- D) $(-0.453\text{ m}, -0.767\text{ m})$
- E) $(-0.670\text{ m}, -0.767\text{ m})$

Q13.

Figure 7 shows a plot of the angular velocity versus time for a disk rotating about a fixed axis through its center. Rank the time intervals according to the magnitude of the angular acceleration, greatest first.

Fig#



- A) c
- B) a
- C) b
- D) d
- E) All tie

Q14.


A wheel is rotating with a constant angular acceleration of -2.0 rad/s^2 . In the first 4.0 seconds, it makes 8.0 revolutions. What is the total number of revolutions (starting from $t = 0$) will it make before stopping?

- A) 11
- B) 16
- C) 19
- D) 22
- E) 14

Q15.

Figure 8 shows a disk with a moment of inertia $I = 10.0 \text{ kg}\cdot\text{m}^2$ about an axis passing through its center. Two strings are wrapped around different parts of the disk which have radii $R_1 = 40.0 \text{ cm}$ and $R_2 = 25.0 \text{ cm}$. Find the magnitude of the angular acceleration of the disk if the tensions are $T_1 = 5.0 \text{ N}$ and $T_2 = 15 \text{ N}$.

Fig#




- A) 0.18 rad/s^2
 - B) 10 rad/s^2
 - C) 0.40 rad/s^2
 - D) 0.25 rad/s^2
 - E) 1.2 rad/s^2
-

Q16.

A meter stick is held vertically with one end pivoted on the floor. It is then allowed to fall as shown in **Figure 9**. Find the speed of the other end just before it hits the floor.

Fig#




- A) 5.4 m/s
 - B) 2.0 m/s
 - C) 7.7 m/s
 - D) 3.3 m/s
 - E) 12 m/s
-

Q17.

Figure 10 shows a uniform rod of length 0.6 m and mass 1.0 kg, rotating in the plane of the figure about an axis through one end. When it is at its lowest point, it collides with a stationary 0.2 kg object that sticks to the end of the rod. If the rod's angular speed just before collision is 2.4 rad/s, then what is its angular speed just after the collision?

Fig#



- A) 1.5 rad/s
- B) 0.52 rad/s
- C) 2.1 rad/s
- D) 1.2 rad/s
- E) 1.9 rad/s

Q18.

At $t = 0$, a 2.0 kg particle with velocity $\mathbf{v} = (5.0 \mathbf{i} + 3.0 \mathbf{j})$ m/s is at the origin. It is pulled by a 6.0 N force in the negative y direction. What is the torque (in units of N.m) about the origin at $t = 3.0$ s?

- A) $-90 \mathbf{k}$
 - B) $-21 \mathbf{k}$
 - C) $88 \mathbf{k}$
 - D) 0
 - E) $62 \mathbf{k}$
-

Q19.


Two wheels A and B of the same radius and mass start rolling from rest, down the same incline (without slipping) from the same initial height. The difference between the two wheels is that wheel A has more mass near the rim while wheel B has more mass near the center. When they reach the bottom, which one of the following statements is TRUE?

- A) Wheel B rolls down faster than wheel A
- B) Wheel A rolls down faster than wheel B
- C) Both wheels roll at the same speed
- D) The change in the potential energy of wheel A at the bottom of the incline is greater than that of wheel B
- E) The wheels will have the same kinetic energy at the bottom of the incline

Q20.

Figure 11 shows a disk of mass = 2.0 kg rolling up an incline ($\theta = 20^\circ$) starting with an initial (total) kinetic energy of 88 J. How far does the disk travel along the incline before stopping momentarily?

Fig#



- A) 13 m
 - B) 4.5 m
 - C) 4.8 m
 - D) 1.7 m
 - E) 2.5 m
-