

Q1.

A constant force $\vec{F} = (7.0\hat{i} - 2.0\hat{j})N$ acts on a 2.0 kg block, initially at rest, on a frictionless horizontal surface. If the force causes the block to be displaced by $\vec{d} = (4.0\hat{i} + 6.0\hat{j})m$, find the block's final speed.

- A) 4.0 m/s
- B) 5.0 m/s
- C) 0 m/s
- D) 3.0 m/s
- E) 2.0 m/s


Q2.

A 15.0 N force with a fixed direction does work on a particle as the particle moves through the three-dimensional displacement $\vec{d} = (2.00\hat{i} - 4.00\hat{j} + 3.00\hat{k})m$. What is the angle between the force and the displacement if the change in the particle's kinetic energy is 50.0 J.

- A) 51.8°
- B) 62.3°
- C) 43.9°
- D) 69.1°
- E) 37.2°

Q3.

The only force acting on a 2.0 kg body as the body moves along an x axis varies as shown in **Figure 1**. The scale of the figure's vertical axis is set by $F_s = 5.0$ N. The speed of the body at $x = 0$ is 5.0 m/s. At what value of x will the body have a kinetic energy of 15 J?



- A) 4.0 m
- B) 1.0 m
- C) 3.0 m
- D) 5.0 m
- E) 2.0 m


Q4.

The loaded cab of an elevator has a mass of 3.0×10^3 kg and moves 150 m upward in 10 s at constant speed. At what average rate does the force from the cable do work on the cab? [Ignore air resistance]

- A) 4.4×10^5 W
- B) 2.0×10^5 W
- C) 2.7×10^5 W
- D) 2.7×10^3 W
- E) 0 W

Q5.

In **Figure 2**, a 6.00 kg block is projected up a frictionless plane at a speed of 4.00 m/s. The plane is inclined at an angle $\theta = 30.0^\circ$ with the horizontal. How far up along the inclined plane does the block move before coming to stop? [Ignore air resistance]



- A) 1.63 m
- B) 0.850 m
- C) 4.00 m
- D) 1.00 m
- E) 1.05 m


Q6.

A man pushes a block up an incline at a constant speed. As the block moves up the incline, only one statement is true:

- A) Its kinetic energy remains constant and the potential energy of block-earth system increases
- B) Its kinetic energy and the potential energy of block-earth system both increase
- C) Its kinetic energy increases and the potential energy of block-earth system remains constant
- D) Its kinetic energy decreases and the potential energy of block-earth system remains constant
- E) Its kinetic energy decreases and the potential energy of block-earth system decreases

Q7.


As shown in **Figure 3**, a 3.50 kg block is accelerated from rest by compressing against a spring by a distance x . The other end of the spring is fixed to the wall and the spring has a spring constant of 134 N/m. The block leaves the spring and then travels over a rough horizontal floor with a coefficient of kinetic friction $\mu_k = 0.250$. The frictional force stops the block in distance $D = 7.80$ m. Find the distance x ?



- A) 1.00 m
- B) 0.460 m
- C) 1.20 m
- D) 0.750 m
- E) 2.30 m

Q8.

A 80 kg skier starts from rest at height $H = 25$ m above the end of a frictionless ski-jump ramp and leaves the ramp at angle $\theta = 30^\circ$, as shown in **Figure 4**. What is the maximum height h of his jump above the end of the ramp? Neglect the effects of air resistance.



- A) 6.3 m
- B) 5.6 m
- C) 7.1 m
- D) 4.3 m
- E) 8.1 m


Q9.

A completely inelastic collision occurs between two balls of clay that move directly toward each other along a vertical axis. Just before the collision, one ball, of mass 5.0 kg, is moving upward at 20 m/s and the other ball, of mass 2.0 kg, is moving downward at 15 m/s. After the collision the two balls stick together and move. How high do the combined two balls rise above the collision point? (Neglect air resistance.)

- A) 5.1 m
- B) 2.6 m
- C) 7.3 m
- D) 3.7 m
- E) 4.5 m

Q10.

A 3.00 kg ball strikes a wall at 10.0 m/s at an angle of 60.0° with the plane of the wall. It bounces off the wall with the same speed and angle (See **Figure 5**). If the ball is in contact with the wall for 0.200 s, what is the magnitude of the average force exerted by the wall on the ball? [ignore air resistance]



- A) 260 N
- B) 150 N
- C) 110 N
- D) 390 N
- E) 0

Q11.

The center of mass of a two-particle system is at the origin. One particle has a mass of 2.0 kg and is located at $(x = 3.0 \text{ m}, y = 0.0 \text{ m})$. What is the location of the second particle if it has a mass of 4.0 kg?

- A) $(x = -1.5 \text{ m}, y = 0.0 \text{ m})$
- B) $(x = +1.5 \text{ m}, y = -1.5 \text{ m})$
- C) $(x = -1.5 \text{ m}, y = +1.5 \text{ m})$
- D) $(x = 0.0 \text{ m}, y = -1.5 \text{ m})$
- E) $(x = +1.5 \text{ m}, y = 0.0 \text{ m})$

Q12.

Two metallic solid spheres approach each other head-on with the same speed of 3.00 m/s and collide elastically. After the collision, one of the spheres, whose mass is 600 g, remains at rest. What is the speed of the two-sphere center of mass?

- A) 1.50 m/s
- B) 3.55 m/s
- C) 1.00 m/s
- D) 2.75 m/s
- E) 4.53 m/s


Q13.

Starting from rest, a disk takes 10 rev. to reach an angular velocity ω at constant angular acceleration. How many more revolutions are required by the disk to reach an angular velocity of 2ω ?

- A) 30 rev.
- B) 10 rev
- C) 20 rev.
- D) 40 rev
- E) 50 rev.

Q14.


A thin uniform rod (mass = 5.0 kg and length = 6.0 m) rotates freely in a vertical plane about a horizontal axis that is perpendicular to the rod and passing through point A, as shown in **Figure 6**. Point A is at distance $d = 1.5$ m from the end of the rod. The kinetic energy of the rod as its other end B passes through the lowest vertical position is 25 J. What is the (linear) speed of the end B as the rod passes through the lowest vertical position?



- A) 6.2 m/s
- B) 2.6 m/s
- C) 3.9 m/s
- D) 7.9 m/s
- E) 8.1 m/s

Q15.

A 45 N block is suspended by a massless light string from a 2.0 kg pulley, as shown in **Figure 7**. The block is released from rest and falls to the floor below as the pulley rotates. The pulley may be considered a solid disk of radius 1.5 m. What is the magnitude of acceleration \vec{a} of the block? [Ignore air resistance]



- A) 8.1 m/s^2
- B) 15 m/s^2
- C) 9.4 m/s^2
- D) 5.5 m/s^2
- E) 7.3 m/s^2

Q16.

A 2.0 kg solid cylinder of radius 0.50 m rotates about its cylindrical axis at a constant angular velocity of 40 rad/s. What average power is required to bring the cylinder to rest in 10 s?

- A) 20 W
- B) 10 W
- C) 50 W
- D) 30 W
- E) 40 W

Q17.

A uniform solid sphere rolls without slipping along a horizontal surface. What percentage of its total kinetic energy is rotational kinetic energy?

- A) 29 %
- B) 50 %
- C) 12 %
- D) 75 %
- E) 33 %

Q18.

Force $\vec{F} = (3\hat{i} + \hat{j})N$ is acting on a particle with position vector $\vec{r} = (2\hat{i} + 4\hat{j})m$. What is resulting torque on the particle about a point ($x = 0$ m, $y = 6$ m)?

- A) $8\hat{k}$ N.m
- B) $-8\hat{k}$ N.m
- C) $5\hat{k}$ N.m
- D) $-5\hat{k}$ N.m
- E) $3\hat{k}$ N.m


Q19.

A point object with 1.40 kg mass and position ($x = 2.00$ m, $y = 3.10$ m) moves at 4.62 m/s at an angle 45.0° north of east. What is magnitude of the object's angular momentum about the origin.

- A) 5.00 kg.m²/s
- B) 3.60 kg.m²/s
- C) 6.00 kg.m²/s
- D) 7.00 kg.m²/s
- E) 2.90 kg.m²/s

Q20.

Two uniform spheres A and B of equal masses, attached to two ends of a massless rod, can rotate about a frictionless pivot at the point P in horizontal plane on a tabletop. **Figure 8** shows top view of the rotating system. The system is made to rotate clockwise on the tabletop with angular speed ω . Sphere A collides with and sticks to another sphere C (with equal mass) that is at rest on the tabletop. What is the angular speed of the system immediately after the collision?



- A) 0.56ω
- B) 0.82ω
- C) 0.60ω
- D) ω
- E) 0.