


Q1.

Figure 1 gives the x component F_x of a single force that acts on a particle. If the particle begins at rest at $x = 0$, what is its coordinate when it has its greatest kinetic energy?



- A) 3 m
- B) 1 m
- C) 6 m
- D) 5 m
- E) 8 m


Q2.

A 100 kg block is pulled at a constant speed of 5.0 m/s across a horizontal floor by an applied force of 122 N directed 37° above the horizontal. What is the rate at which the force does work on the block?

- A) 4.9×10^2 W
- B) 3.7×10^2 W
- C) 5.6×10^2 W
- D) 1.8×10^3 W
- E) 2.4×10^1 W

Q3.

A 0.80 kg block is dropped onto a relaxed vertical spring that has a spring constant of $k = 250$ N/m as shown in **Figure 2**. The block compresses the spring 0.12 m before momentarily stopping. Find the maximum speed of the block just before it hits the spring. (Assume that friction and air resistance are negligible.)



- A) 1.5 m/s
- B) 2.1 m/s
- C) 3.2 m/s
- D) 4.6 m/s
- E) 5.0 m/s

Q4.

A single force \vec{F} acts on a 0.40-kg particle and changes its velocity from $\vec{v}_i = (4.0\hat{i} - 3.0\hat{j})$ m/s at time t_i to $\vec{v}_f = (5.0\hat{i} + 3.0\hat{j})$ m/s at time t_f . What is the work done by \vec{F} on the particle during this interval of time?

- A) 1.8 J
- B) 0.14 J
- C) 1.2 J
- D) zero
- E) 5.0 J


Q5.

At time $t = 0$, a 1.0 kg ball is thrown from the top of a 100 m tall tower with initial velocity $\vec{v}_0 = (16\hat{i} + 24\hat{j})$ m/s. At what height from the ground will the kinetic energy of the ball be three times its initial kinetic energy? (Ignore the air resistance)?

- A) 15 m
- B) 10 m
- C) 20 m
- D) 25 m
- E) 40 m

Q6.

A block with mass $m = 2.00$ kg is placed against a spring on a rough incline with angle $\theta = 30.0^\circ$ and coefficient of kinetic friction $\mu_k = 0.215$ as shown in **Figure 3** (The block is not attached to the spring). The spring, which is compressed 20.0 cm from its relaxed position, is then released from rest and the block travels distance $l = 1.20$ m from the release point on the incline before coming to rest. Find the value of spring constant k of the spring.



- A) 807 N/m
- B) 578 N/m
- C) 256 N/m
- D) 980 N/m
- E) 663 N/m

Q7.

If only conservative forces are acting on a body then the work done by conservative forces

- A) does not change the total mechanical energy.
 - B) does not change the potential energy.
 - C) does not change the kinetic energy.
 - D) is always equal to zero.
 - E) is always negative.
-

Q8.

An 18-kg object is released from rest and moves vertically downward from a height of 80 m above the ground. It reaches the ground with a speed of 15 m/s. How much work was done by the non-conservative forces on the object?

- A) - 12 kJ
 - B) - 16 kJ
 - C) + 12 kJ
 - D) + 16 kJ
 - E) - 14 kJ
-

Q9.

A stone is dropped at time $t = 0$. A second stone, with twice the mass of the first, is dropped from the same point at $t = 0.10$ s. How far below the release point is the center of mass of the two stones at $t = 0.30$ s? Ignore air resistance. (Both stones are dropped from rest and none of the stones has reached the ground.)

- A) 0.28 m
 - B) 0.12 m
 - C) 0.45 m
 - D) 0.31 m
 - E) 0.63 m
-

Q10.

A 2.4-kg ball that is falling vertically downward hits a horizontal floor with a speed of 2.5 m/s and rebounds with a speed of 1.5 m/s. What is the magnitude of the impulse exerted on the ball by the floor?

- A) 9.6 N.s
 - B) 2.4 N.s
 - C) 3.5 N.s
 - D) 6.7 N.s
 - E) 7.1 N.s
-


Q11.

A cart, with mass 340 g and moving on a horizontal frictionless surface with an initial speed of 1.2 m/s, undergoes an elastic collision with an initially stationary cart of unknown mass. After the collision, the first cart continues in its original direction at 0.66 m/s. What is the mass of the second cart?

- A) 0.099 kg
 - B) 0.061 kg
 - C) 0.036 kg
 - D) 0.018 kg
 - E) 0.075 kg
-

Q12.

A 4.0 kg mass, moving with constant speed v , explodes at point O into two equal parts, as shown in **Figure 4**. The first part moves with speed 3.0 m/s due north, and the second part moves with speed 5.0 m/s, 30° north of east. Find the value of v .



- A) 3.5 m/s
- B) 8.0 m/s
- C) 5.0 m/s
- D) 2.0 m/s
- E) 4.5 m/s


Q13.

A rotating wheel requires 3.00 s to rotate through 37.0 revolutions. Its angular speed at the end of the 3.00 s interval is 98.0 rad/s. What is the constant angular acceleration of the wheel?

- A) 13.7 rad/s²
- B) 10.5 rad/s²
- C) 11.2 rad/s²
- D) 17.1 rad/s²
- E) 29.3 rad/s²

Q14.

Find the net torque on the wheel in **Figure 5** about the axle through O if $a = 10.0$ cm and $b = 25.0$ cm.



- A) -3.55 N.m
- B) -1.27 N.m
- C) $+1.27$ N.m
- D) $+3.55$ N.m
- E) -7.16 N.m


Q15.

A 32.0 kg wheel, essentially a thin hoop with radius 1.20 m, is rotating about its axis at 280 rev/min. It must be brought to a stop in 15.0 s. What is the magnitude of the required average power to stop it?

- A) 1.32×10^3 W
- B) 2.53×10^3 W
- C) 6.14×10^3 W
- D) 3.51×10^3 W
- E) 4.96×10^3 W

Q16.


A mass ($M_1 = 5.0$ kg) is connected by a massless cord to another mass ($M_2 = 4.0$ kg) which slides on a horizontal frictionless surface, as shown in **Figure 6**. The pulley (radius = 0.20 m) rotates about a frictionless axle. If the acceleration of M_2 is 3.5 m/s², what is the rotational inertia of the pulley?



- A) 0.20 kg.m²
- B) 0.50 kg.m²
- C) 0.10 kg.m²
- D) 0.35 kg.m²
- E) 0.75 kg.m²

Q17.


A uniform solid sphere of radius 0.10 m started to roll up without slipping with a center of mass speed of 2.0 m/s from the bottom of a ramp (point A in **Figure 7**) that is inclined at an angle $\theta = 10^\circ$. Find the maximum distance (d) travelled by the ball before it comes to rest.



- A) 1.6 m
- B) 2.8 m
- C) 3.9 m
- D) 4.1 m
- E) 6.3 m

Q18.

Figure 8 gives the angular momentum magnitude L of a wheel versus time t . Rank the four lettered time intervals according to the magnitude of the torque acting on the wheel, **greatest first**.



- A) D, B, (A and C) tie
- B) B, (A and C) tie, D
- C) D, (A and C) tie, B
- D) (A and C) tie, B, D
- E) B, D, (A and C) tie


Q19.

A 2.0 kg particle-like object moves in an xy -plane with velocity components $v_x = 20$ m/s and $v_y = 60$ m/s as it passes through the point with (x, y) coordinates of $(3.0, -4.0)$ m. At this time, what are the magnitude (in SI units) and direction of its angular momentum relative to the point located at $(-2.0, -2.0)$ m?

- A) $+680 \hat{k}$
- B) $-680 \hat{k}$
- C) Zero
- D) $+540 \hat{k}$
- E) $-540 \hat{k}$

Q20.

A 50 g bullet is fired horizontally at one end of a 0.60 m long uniform rod of mass 0.50 kg, which is originally at rest and is pivoted at another end at point A in a vertical plane, as shown in **Figure 9**. If the angular speed of the system (bullet + rod assumed to stick together) about A just after impact is 4.5 rad/s, what is the bullet's speed just before impact?



- A) 12 m/s
 - B) 10 m/s
 - C) 14 m/s
 - D) 17 m/s
 - E) 20 m/s
-