


Q1.

Figure 1 shows a graph of the acceleration versus the displacement of a particle moving in one dimension. The particle is at rest at $x = 0$. What is the coordinate x at which the particle has the maximum kinetic energy?



- A) 3
- B) 1
- C) 5
- D) 7
- E) 8

Q2.

What is the work done by the force $\vec{F} = (2.0\hat{i} + 4.0\hat{j} + 9.0\hat{k})$ N that acts on a 3.0 kg object and moves from an initial position $\vec{r}_1 = (-4.1\hat{i} + 3.3\hat{j} + 5.2\hat{k})$ m to a final position $\vec{r}_2 = (2.7\hat{i} - 2.9\hat{j} + 5.5\hat{k})$ m.

- A) -8.5 J
- B) -12 J
- C) 5.2 J
- D) 12 J
- E) 4.7 J

Q3.

A 25 kg block, which is initially at rest, is pulled across a horizontal rough surface (coefficient of kinetic friction $\mu_k = 0.3$) by a force of 80 N directed 30° above the horizontal. What is the final speed of the block after it has moved a distance $d = 10$ m?

- A) 2.5 m/s
- B) 1.8 m/s
- C) 3.3 m/s
- D) 4.0 m/s
- E) 3.8 m/s


Q4.

A worker uses a motor to raise a 2700-kg block a vertical distance of 3.2 m. If the task is to be achieved in 5.0 minutes, what is the minimum (average) power requirement for the motor?

- A) 280 W
- B) 320 W
- C) 440 W
- D) 510 W
- E) 560 W

Q5.


A block of mass $m = 3.5$ kg slides from rest down a frictionless incline of angle $\theta = 30^\circ$. See **Figure 2**. After sliding a distance d along the incline, it compresses a relaxed spring of spring constant 430 N/m. The block momentarily stops after compressing the spring by 20 cm. What is the distance d ?



- A) 30 cm
- B) 40 cm
- C) 50 cm
- D) 55 cm
- E) 60 cm

Q6.


A box can slide along a track with elevated curved ends and a flat central part, as shown in **Figure 3**. The flat part has length $2L$. The curved portions of the track are frictionless, but for the flat part the coefficient of kinetic friction is $\mu_k = 0.25$. The box is released from rest at point A on the left curved portion, which is at a height L . Find the maximum height that the box will reach at the right curved portion.



- A) $L/2$
 B) L
 C) $3L/2$
 D) $2L/3$
 E) $L/3$

Q7.


A pendulum consists of a small mass $m = 0.15$ kg attached to a massless rod with length 2.0 m. At $\theta_0 = 15^\circ$, the mass has a speed of 2.0 m/s (see **Figure 4**). Find the maximum angle θ_{\max} that the pendulum will make with the vertical. (Ignore air resistance)



- A) 30°
 B) 37°
 C) 27°
 D) 45°
 E) 60°

Q8.

A ball of mass 0.15 kg and with initial speed 2.0 m/s, collides elastically with the floor at $\theta = 30^\circ$ with the vertical and bounces back with the same speed and the same angle (see **Figure 5**). What is the impulse from the ball on the floor?



- A) 0.52 N.s along the negative y-axis
 - B) 0.52 N.s along the positive y-axis
 - C) 0.30 N.s along the positive x-axis
 - D) 0.30 N.s along the negative x-axis
 - E) 0
-

Q9.

Figure 6, shows three blocks A, B, and C, of masses $3M$, M and $2M$ respectively, on a frictionless surface. Blocks B and C are initially at rest. Block A is initially moving towards block B at a speed v and undergoes an elastic collision with block B. Block B moves to the right and undergoes a completely inelastic collision with block C. What is the speed of block C immediately after the collision?



- A) $v/2$
 - B) $v/3$
 - C) $3v/4$
 - D) $v/4$
 - E) $2v/3$
-

Q10.

Object A of mass m is moving along the x -axis with an initial speed of 5.0 m/s. It then collides with another object B of mass $4m$, which is initially at rest. After the collision, object A moves with a speed of 2.5 m/s in a direction 60° to its original direction of motion. If the collision is inelastic, determine the direction of travel of object B with respect to the x -axis.

- A) 30°
- B) 37°
- C) 60°
- D) 45°
- E) 53°

Q11.

A stationary block lying on a frictionless floor explodes into three pieces that slide across the floor. Which one of the following quantities **will NOT be zero** after the explosion?

- A) The total kinetic energy
- B) The velocity of the center of mass
- C) The acceleration of the center of mass
- D) The total linear momentum
- E) The displacement of the center of mass


Q12.

A car and a truck, initially at rest at a traffic light, start accelerating in the same direction when the traffic light turns green. The acceleration of the car is 15.0 m/s^2 and the acceleration of the truck is 5.0 m/s^2 . If the mass of the truck is three times the mass of the car, then what is the magnitude of the velocity of their center of mass after 4.0 s ?

- A) 30 m/s
- B) 20 m/s
- C) 40 m/s
- D) 25 m/s
- E) 35 m/s

Q13.

Four equal masses are arranged at the corners of a rectangle of width L and length $2L$. The masses are connected by rigid, massless rods. The system can rotate about any one of the axes 1, 2, or 3 shown in **Figure 7**. Rank the axes according to the value of the rotational inertia about them, **greatest to least**.



- A) 2, 1, 3
- B) 2, 3, 1
- C) 1, 2, 3
- D) 1, 3, 2
- E) 3, 2, 1

Q14.

A rotating ring has a radius $R = 22$ m. The angular position of a reference line on the ring is given by: $\theta(t) = 1.5t^3 - 4.0t^2$. Find the ratio of the magnitude of the tangential acceleration to the magnitude of the radial acceleration at $t = 2.0$ s for a point on the rim of the ring.

- A) 2.5
- B) 3.0
- C) 4.0
- D) 5.5
- E) 4.4


Q15.

A wheel is initially rotating at 10 rad/s and has a constant angular acceleration. After 9.0 s, it has rotated through 24 revolutions. What is the magnitude of its angular acceleration?

- A) 1.5 rad/s²
- B) 3.5 rad/s²
- C) 4.4 rad/s²
- D) 5.3 rad/s²
- E) 5.9 rad/s²

Q16.

A yo-yo shaped device mounted on a horizontal frictionless axis is used to lift a 25 kg box as shown in **Figure 8**. The box is suspended by a rope wrapped around the axle whose radius r is 0.30 m. The outer radius R of the device is 0.6 m. When a constant horizontal force F of magnitude 140 N is applied to the rope at the outer radius of the device, the box has an upward acceleration of magnitude 0.80 m/s². What is the rotational inertia of the device about its axis of rotation?



- A) 1.7 kg m²
- B) 2.6 kg m²

- C) 4.2 kg m^2
D) 5.0 kg m^2
E) 3.2 kg m^2

Q17.

A 0.50-kg object moves in a horizontal circular track with radius of 2.0 m . An external force of 4.0 N always tangent to the track causes the object to speed up as it goes around. What is the work done by the external force as the object makes one revolution?

- A) 50 J
B) 60 J
C) 70 J
D) 80 J
E) 0 J

Q18.


A uniform solid sphere rolls smoothly without slipping with center of mass velocity of 2.5 m/s along a horizontal floor, then up a ramp inclined at 20° . What is the maximum distance traveled by the center of mass of the sphere along the ramp?

- A) 1.3 m
B) 1.0 m
C) 1.7 m
D) 2.1 m
E) 2.4 m

Q19.

Figure 9 is a plot of the angular momentum of a wheel versus time. At which time interval (A, B, C or D) is the magnitude of the torque acting on the wheel maximum?


L ($\text{kg m}^2/\text{s}$)



- A) D only
B) A only
C) B only
D) C only
E) A and C

Q20.

The uniform thin rod in **Figure 10** (mass $M = 3m_o$ and length $L = 1.5$ m) can rotate about a vertical axis through one end (O). Initially, it is held horizontally. When released, it swings through its lowest position and collides with a stationary block of mass m_o that sticks to the end of the rod. What is the angular speed (in rad/s) of the rod-block system immediately after the collision? Take $m_o = 0.50$ kg. (Ignore all forms of frictions)



- A) 2.2
- B) 3.2
- C) 1.5
- D) 3.6
- E) 4.0