

Q1.

A ball with a weight of 4.5 N is thrown at an angle of 30° above the horizontal with an initial speed of 10 m/s. Neglecting air resistance, at its highest point, the net force on the ball is:

- A) 4.5 N, vertically downward
- B) Zero
- C) 1.5 N, 30° below horizontal
- D) 9.8 N, vertically downward
- E) 9.8 N, 30° below horizontal

Q2.

A ball is suspended by a string from the ceiling of a car. The car moves horizontally with a constant acceleration of 2.5 m/s^2 with respect to the ground. The ball is at rest with respect to the car. What angle does the string make with the vertical?

- A) 14°
- B) Zero
- C) 25°
- D) 45°
- E) 90°


Q3.

A 5.0 kg block is lowered with a downward acceleration of 2.8 m/s^2 by means of a rope. Find the force exerted by the block on the rope.

- A) 35 N, downward
- B) 14 N, downward
- C) 35 N, upward
- D) 14 N, upward
- E) 49 N, upward

Q4.


Two blocks of masses $m_1 = 2.0 \text{ kg}$ and $m_2 = 1.0 \text{ kg}$ are in contact with each other on a frictionless surface as shown in **Figure 1**. A horizontal force \vec{F} of magnitude 3.0 N is applied to the block of mass m_2 as shown. Find the magnitude of the force that m_1 exerts on m_2 .



- A) 2.0 N
- B) 9.8 N
- C) 1.0 N
- D) 6.0 N
- E) 3.0 N

Q5.


A pickup truck carries a box on its back as shown in **Figure 2**. The truck starts from rest. When the truck attains an acceleration of 2.00 m/s^2 , the box starts to slip towards the end of the truck. The coefficient of friction between the box and the truck's back surface is ?



- A) $\mu_s = 0.20$
- B) $\mu_s = 0.15$
- C) $\mu_s = 0.37$
- D) $\mu_s = 0.11$
- E) $\mu_s = 0.30$

Q6.

A block is sliding down a rough inclined plane that makes an angle of 30° with the horizontal. When a horizontal force \vec{F} of magnitude 10 N is applied to the block, as shown in **Figure 3**, the block slides down at a constant speed. If the coefficient of kinetic friction between the block and the inclined plane is 0.30, find the mass of the block.



- A) 4.3 kg
- B) 1.5 kg
- C) 4.7 kg
- D) 2.0 kg
- E) 3.9 kg

Q7.

A car of mass 500 kg can go around a banked circular road of radius 60 m at the maximum speed of 72 km/h without slipping. Find the normal force on the car from the banked surface (Ignore the friction force from the road).

- A) 5.9 kN
- B) 6.4 kN
- C) 39 kN
- D) 1.9 kN
- E) 79 kN

Q8.

A block is dropped from a high tower and is falling freely under the influence of gravity. Which one of the following statements is **true**? (Ignore air resistance).

- A) The kinetic energy increases by equal amounts over equal distances.
- B) As the block falls, the net work done by all of the forces acting on the block is zero joules.
- C) The kinetic energy of the block increases by equal amounts in equal times.
- D) The potential energy of the block decreases by equal amounts in equal times.
- E) The total energy of the block increases by equal amounts over equal distances.


Q9.

A massless spring hangs from the ceiling. How much does its potential energy increase if a 30 kg mass is attached to it? The spring constant is 4800 N/m.

- A) 9.0 J
- B) 1.9 J
- C) 5.3 J
- D) 12 J
- E) 15 J

Q10.

A 10 kg block is sent up a frictionless ramp along which an x axis extends upward. **Figure 4** gives the kinetic energy of the block as a function of position x ; the scale of the figure's vertical axis is set by $K_s = 80$ J. What is the angle of the inclination of the ramp with respect to the horizontal surface?



- A) 24°
- B) 31°
- C) 19°
- D) 33°
- E) 11°

Q11.

Starting from rest, an elevator with a mass of 1.00×10^3 kg moves 100 m vertically upward in 50.0 s. At what average rate does the force from the cable do work on the elevator?

- A) 19.8 kW
- B) 11.1 kW
- C) 29.7 kW
- D) 31.3 kW
- E) 15.6 kW

Q12.

Two balls are launched from the same spring-loaded cannon with the spring compressed the same distance each time. Ball A has a 40 kg mass and ball B has a 60 kg mass. The relation between their speeds at the instant of launch is:

- A) $v_A = \sqrt{(3/2)} v_B$
 B) $v_A = v_B$
 C) $v_A = (3/2) v_B$
 D) $v_B = \sqrt{(3/2)} v_A$
 E) $v_B = (3/2) v_A$


Q13.

In a simple pendulum a 2.00 kg mass is attached to a 2.00 m long massless string. The mass has a speed of 3.00 m/s when the string makes an angle of 30.0° with the vertical. What is the speed of the mass when the string makes an angle of 45° with the vertical?

- A) 1.66 m/s
 B) 2.64 m/s
 C) 5.22 m/s
 D) 26.6 m/s
 E) 0.720 m/s

Q14.


Figure 5 shows a 10.0 kg stone resting on a spring. The spring is compressed 10.0 cm by the stone. The stone is further pushed down an additional 40.0 cm and then released. The stone rises vertically to a maximum height of 1.00 m from its release point. What is the magnitude of the work done by non-conservative forces (air resistance) on the stone during its flight to the maximum height?



- A) 24.5 J
 B) 14.7 J
 C) 34.9 J
 D) 9.25 J
 E) 41.5 J

Q15.

Starting from rest, a 2.00 kg block slides downward inside a frictionless circular hoop of radius $R = 0.50$ m, as shown in **Figure 6**. What is the magnitude of the normal force exerted on the block by the hoop when the block reaches the bottom of the hoop?



- A) 58.8 N
 - B) 88.5 N
 - C) 30.1 N
 - D) 72.4 N
 - E) 31.7 N
-
-