

**Q1.** A light body and a heavy body have equal linear momenta. The one having the larger kinetic energy is:


- A) The light body.
- B) The heavy body.
- C) Neither; they will have the same kinetic energy.
- D) Dependent on the system of units used.
- E) Not determinable without data on the ratio of the masses.

**Q2.** Two particles of masses 3.00 kg and 5.00 kg are moving with velocities of:  $(-3.00\hat{i} + 4.00\hat{j})$  m/s and  $(2.00\hat{i} + 3.00\hat{j})$  m/s, respectively. They collide completely inelastically. Find the velocity of the center of mass after collision.

- A)  $(0.125\hat{i} + 3.38\hat{j})$  m/s
- B)  $(3.23\hat{i} + 4.32\hat{j})$  m/s
- C)  $(5.13\hat{i} + 1.34\hat{j})$  m/s
- D)  $(-9.00\hat{i} + 1.34\hat{j})$  m/s
- E)  $(10.0\hat{i} + 15.0\hat{j})$  m/s

**Q3.** A uniform and thin rectangular piece of wood of width 20 cm and length 50 cm has a mass of 2.0 kg. Two point masses 3.0 kg and 5.0 kg are attached to it at points A and B, respectively (see **Figure 1**). Find the  $x$  and  $y$  coordinates, respectively, of the center of mass of the system relative to the origin.

- A) (35 cm, 8.0 cm)
- B) (25 cm, 10 cm)
- C) (30 cm, 20 cm)
- D) (50 cm, 10 cm)
- E) (20 cm, 20 cm)




**Q4.** A body, initially at rest, suddenly explodes into two fragments of masses 0.100 kg and 0.500 kg. The 0.500 kg fragment moves in the positive  $x$  direction. Of the energy released in the explosion, ONLY  $9.60 \times 10^3$  J were converted into kinetic energy of the two fragments. Calculate the speed of 0.100 kg and 0.500 kg fragments, respectively.

- A) 400 m/s, 80.0 m/s
- B) 250 m/s, 150 m/s
- C) 300 m/s, 300 m/s
- D) 300 m/s, 500 m/s
- E) 500 m/s, 300 m/s

**Q5.** A weight  $W = 100 \text{ N}$  is hung from two ropes as shown in **Figure 2**. Find the magnitude of the tension in the horizontal rope.

- A) 173 N
- B) 410 N
- C) 650 N
- D) 321 N
- E) 258 N


Figure # 2



**Q6.** Two antiparallel forces of equal magnitudes  $F_1 = F_2 = 8.00 \text{ N}$  are applied to a rod as shown **Figure 3**. Find the distance  $l$  between the forces if the magnitude of the net torque due to these two forces about the end  $O$  is  $6.40 \text{ N}\cdot\text{m}$ .

- A) 0.800 m
- B) 0.540 m
- C) 0.235 m
- D) 0.458 m
- E) 0.995 m


Figure # 3



**Q7.** A particle of mass  $m$  moving in the positive  $x$  direction with speed  $u$  collides with a particle of mass  $2m$  at rest. After collision, the particle of mass  $m$  scatters with speed  $u/4$  in the positive  $y$  direction and the particle of mass  $2m$  moves with speed  $v$  making an angle  $\theta$  with the positive  $x$  direction (see **Figure 4**). Find the angle  $\theta$ .

- A)  $14^\circ$
- B)  $25^\circ$
- C)  $35^\circ$
- D)  $45^\circ$
- E)  $55^\circ$

Figure # 4



**Q8.** A force of  $5.00 \times 10^3$  N is applied outwardly perpendicular to one end of a 5.00 m long cylindrical rod with a radius of 34.0 cm and a Young's modulus of  $1.25 \times 10^8$  N/m<sup>2</sup>, while the other end is tightly fixed to the wall. Find the elongation of the rod.

- A) 0.551 mm
- B) 0.263 mm
- C) 0.149 mm
- D) 0.348 mm
- E) 0.644 mm

**Q9.** A constant torque of 25.0 N.m is applied to a disk that has a rotational inertia of 0.130 kg.m<sup>2</sup> about an axis passing through its center. Find the angular speed after the disk has made 15.0 revolutions starting from rest.


- A) 190 rad/s
- B) 120 rad/s
- C) 155 rad/s
- D) 138 rad/s
- E) 105 rad/s

**Q10.** A turntable rotates with constant  $2.25$  rad /s<sup>2</sup> angular acceleration. After 4.00 s it has rotated through an angle of 60.0 rad. What was the angular speed of the wheel at the beginning of the 4.00 s interval?

- A) 10.5 rad/s
- B) 20.6 rad/s
- C) 15.8 rad/s
- D) 42.6 rad/s
- E) 32.7 rad/s


**Q11.** A thin uniform rod with mass  $M$  and length  $L$  is hinged at one end and connected to the wall, as shown in **Figure 5**. Initially, the rod is held out horizontally then released. Then the magnitude of the rod's angular velocity just before it hits the wall:

- A)  $\sqrt{\frac{3g}{L}}$
- B)  $\sqrt{\frac{2g}{L}}$
- C)  $\sqrt{\frac{7g}{4L}}$
- D)  $\sqrt{\frac{5g}{4L}}$
- E)  $\sqrt{\frac{L}{g}}$



**Q12.** A uniform solid sphere is rolling without slipping along a horizontal surface with a speed of 5.50 m/s when it starts rolling up a ramp that makes an angle of  $30.0^\circ$  with the horizontal. Find the speed of the sphere after it has rolled 3.00 m up the ramp, measured along the surface of the ramp (see **Figure 6**).

- A) 3.04 m/s
- B) 8.02 m/s
- C) 1.91 m/s
- D) 2.16 m/s
- E) 5.37 m/s



**Q13.** The angular momentum of a system remains constant

- A) When no net external torque acts on the system.
- B) When the total kinetic energy is constant.
- C) When no net external force acts on the system.
- D) When the linear momentum and the energy are constant.
- E) All the time since it is a conserved quantity.

**Q14.** An object is rotating with an angular momentum of magnitude  $20.0 \text{ kg}\cdot\text{m}^2/\text{s}$  in the east direction. A torque of magnitude  $10.0 \text{ N}\cdot\text{m}$  in a direction of  $30^\circ$  north of east acts on the object for  $5.00 \text{ s}$ . Find the magnitude of angular momentum at the end of the  $5.00 \text{ s}$  interval.

- A)  $68.1 \text{ kg}\cdot\text{m}^2/\text{s}$
  - B)  $45.0 \text{ kg}\cdot\text{m}^2/\text{s}$
  - C)  $25.0 \text{ kg}\cdot\text{m}^2/\text{s}$
  - D)  $93.3 \text{ kg}\cdot\text{m}^2/\text{s}$
  - E)  $32.0 \text{ kg}\cdot\text{m}^2/\text{s}$
- 

**Q15.** A piece of putty is dropped vertically onto a freely rotating turntable and gets stuck to the turntable. If the rotational inertia of the putty about the center of the turntable is  $0.54$  times that of the turntable about its center. Find the ratio of the final rotational kinetic energy to the initial rotational kinetic energy of the turntable.

- A) 0.65
  - B) 0.18
  - C) 0.46
  - D) 0.28
  - E) 0.87
-