

Q1.

The distance between two successive minima of a transverse wave is 2.76 m. Six crests of the wave pass a given point along the direction of travel every 14.0 s. Find the wave speed.

- A) 0.986 m/s
- B) 0.892 m/s
- C) 0.267 m/s
- D) 1.012 m/s
- E) 0.768 m/s

---

Q2.

The speed of waves on a thin wire is 150 m/s. The density of the material that the wire is made of is  $5000 \text{ kg/m}^3$ . The wire has a 0.500 mm diameter circular cross-section. What is the tension in the wire?

- A) 22.1 N
- B) 76.2 N
- C) 88.4 N
- D) 0.147 N
- E) 63.7 N

---

Q3.


Two identical waves, with amplitude  $A$ , travel simultaneously through the same medium and in the same direction with phase difference  $\phi$ . If the amplitude of the resulting superposition is  $A/2$ , what is the possible value of  $\phi$ ?

- A)  $151^\circ$
  - B)  $45^\circ$
  - C)  $37^\circ$
  - D)  $55^\circ$
  - E)  $91^\circ$
-

Q4.

**Figure1** shows two loudspeakers, (1) and (2), above each other. They are driven by the same source at frequency of 450 Hz. An observer is sitting at point  $O$ , at the same distance from each speaker. What minimum upward vertical distance speaker (1) should be moved to in order to create destructive interference at point  $O$ ? [Note: speed of the sound in air is 343 m/s]

Fig#



- A) 1.43 m
- B) 2.80 m
- C) 1.22 m
- D) 1.01 m
- E) 2.15 m

Q5.

At a distance of 2.0 m from a point source of sound, the sound level is 80 dB. What will be the sound level at a distance of 4.0 m from this source? (Assume that the point source radiates uniformly in all directions)

- A) 74 dB
- B) 83 dB
- C) 62 dB
- D) 16 dB
- E) 12 dB

Q6.

A fixed alarm is emitting sound waves of frequency 520 Hz. You are on a motorcycle, traveling directly away from the alarm. How fast you must be traveling if you detect a frequency of 490 Hz? [Note: Speed of the sound in air = 343 m/s]

- A) 19.8 m/s
- B) 22.2 m/s
- C) 27.1 m/s
- D) 11.9 m/s
- E) 16.3 m/s

Q7.

A 0.90 m long pipe is open at one end but closed at the other end. If it currently resonates with a harmonic of wavelength 0.72 m, what is the wavelength of the next higher harmonic in this pipe?

- A) 0.51 m
- B) 0.33 m
- C) 0.21 m
- D) 0.74 m
- E) 0.88 m

---

Q8.

A point source, of sound waves, radiates uniformly in all directions. At a distance of 20 m from the source the sound level is 51 dB. What is the total power output of the source?

- A)  $6.3 \times 10^{-4}$  Watt
- B)  $9.6 \times 10^{-4}$  Watt
- C)  $1.1 \times 10^{-4}$  Watt
- D)  $9.8 \times 10^{-5}$  Watt
- E)  $4.2 \times 10^{-5}$  Watt

---

Q9.

The coefficient of linear expansion of iron is  $10^{-5} \text{ }^\circ\text{C}^{-1}$ . The volume of an iron cube, having an edge of 5.00 cm, will increase if it is heated from 10.0  $^\circ\text{C}$  to 60.0  $^\circ\text{C}$  by

- A) 0.188 cm<sup>3</sup>
- B) 0.375 cm<sup>3</sup>
- C) 0.225 cm<sup>3</sup>
- D) 0.750 cm<sup>3</sup>
- E) 0.625 cm<sup>3</sup>

---

Q10.


In an insulated container, 250 g of ice at 0  $^\circ\text{C}$  are added to 500 g of water at 18  $^\circ\text{C}$ . What is the final temperature of the system?

- A) 0  $^\circ\text{C}$
- B) 5  $^\circ\text{C}$
- C) 25  $^\circ\text{C}$
- D) 17  $^\circ\text{C}$
- E) 100  $^\circ\text{C}$

Q11.

A system of an ideal gas undergoes the cyclic process shown in the **Figure 2**. Calculate the work done by the system along the path XY.

Fig#




- A) +90 J
- B) -60 J
- C) +60 J
- D) zero
- E) -90 J

Q12.

A bar of gold is in thermal contact with a bar of silver having the same length and area, see **Figure 3**. One end of the connected bars is maintained at  $80.0\text{ }^\circ\text{C}$  and the opposite end is at  $30.0\text{ }^\circ\text{C}$ . When the energy transfer reaches steady state, what is the temperature at the junction? [The thermal conductivity for gold =  $314\frac{\text{W}}{\text{m}\cdot^\circ\text{C}}$  and the thermal conductivity for silver =  $427\frac{\text{W}}{\text{m}\cdot^\circ\text{C}}$ ]

Fig#



- A)  $51.2\text{ }^\circ\text{C}$
- B)  $58.8\text{ }^\circ\text{C}$
- C)  $70.8\text{ }^\circ\text{C}$
- D)  $33.4\text{ }^\circ\text{C}$
- E)  $42.7\text{ }^\circ\text{C}$

Q13.


Two moles of an ideal gas are in a  $6.0 \times 10^{-3} \text{ m}^3$  container at a pressure of  $5.0 \times 10^5 \text{ Pa}$ . Find the average translational kinetic energy of a single molecule.

- A)  $3.7 \times 10^{-21} \text{ J}$
- B)  $7.5 \times 10^{-21} \text{ J}$
- C)  $9.4 \times 10^{-21} \text{ J}$
- D)  $0.22 \times 10^{-21} \text{ J}$
- E)  $5.7 \times 10^{-21} \text{ J}$

Q14.

Which one of the graphs in **Figure 4** best represents the variation of pressure with volume for an isothermal process of an ideal gas?

Fig#



- A) 3
- B) 2
- C) 1
- D) 5
- E) 4

Q15.

One mole of an ideal monatomic gas is initially at 300 K and 1.0 atm. The gas is compressed adiabatically to 2.0 atm. What is the final volume of the gas?

- A)  $0.016 \text{ m}^3$
- B)  $0.037 \text{ m}^3$
- C)  $0.056 \text{ m}^3$
- D)  $0.025 \text{ m}^3$
- E)  $0.012 \text{ m}^3$

Q16.


Find the change in entropy of a 100 g of ice at  $0^\circ\text{C}$  that is isobarically heated slowly to reach  $80^\circ\text{C}$  water. [The heat of fusion for ice  $L_F = 80 \text{ cal/g}$  and the specific heat of water  $c_w = 1 \text{ cal/g.K}$ ].

- A) 55 cal/K
- B) 12 cal/K
- C) 62 cal/K
- D) 35 cal/K
- E) 85 cal/K

Q17.

A sample of an ideal monatomic gas undergoes the reversible process from A to B as displayed in the T-S diagram shown in **Figure 5**. The process is :

Fig#



- A) an isothermal expansion
- B) a free expansion.
- C) an isothermal compression.
- D) a change of phase.
- E) a constant-volume process.

Q18.

Five moles of an ideal monatomic gas are allowed to expand isobarically. The initial volume is  $20.0 \text{ cm}^3$  and the final volume is  $100 \text{ cm}^3$ . Find the change in entropy of the gas.

- A) 168 J/K
- B) 100 J/K
- C) 66.9 J/K
- D) 234 J/K
- E) 33.4 J/K

---

Q19.

A Carnot heat engine absorbs 70.0 kJ as heat and expels 55.0 kJ as heat in each cycle. If the low-temperature reservoir is at  $120 \text{ }^\circ\text{C}$ , find the temperature of the high-temperature reservoir.

- A)  $227 \text{ }^\circ\text{C}$
- B)  $500 \text{ }^\circ\text{C}$
- C)  $153 \text{ }^\circ\text{C}$
- D)  $773 \text{ }^\circ\text{C}$
- E)  $35.9 \text{ }^\circ\text{C}$

---

Q20.

Is it possible to transfer energy as heat from a low-temperature reservoir to a high-temperature reservoir? Choose the right answer with the right explanation.

- A) Yes, this is what a refrigerator does, but work must be done on the refrigerator to make this happen.
  - B) No, this violates the second law of thermodynamics, if no work is being involved.
  - C) No, this violates the zero's law of thermodynamics.
  - D) Yes, this is what a heat engine does, and it can happen without the engine doing work.
  - E) Yes, this is what a refrigerator does, and it can happen without doing work on the refrigerator.
-