

Q1.

A transverse wave traveling on a string is represented by the function  $y(x, t) = y_m \sin[(0.50 \text{ m}^{-1})x - (7.00 \text{ s}^{-1})t]$  where  $x$  is in m and  $t$  is seconds. For which value of the wave amplitude  $y_m$ , the maximum transverse speed  $u_{\text{max}}$  of a particle on the string equals the wave speed  $v$  on the string.

- A) 2.00 m
- B) 1.10 m
- C) 2.57 m
- D) 2.87 m
- E) 3.33 m

Q2.

A guitar string with 0.550 m length and linear density  $\mu = 1.150 \text{ g/m}$  is supposed to have a required fundamental frequency 256 Hz. It currently has a fundamental frequency 248 Hz. What change in tension in the string is required to bring this guitar string into tune with the required frequency?

- A) 5.61 N
- B) 1.23 N
- C) 3.55 N
- D) 4.22 N
- E) 2.79 N

Q3.

A transverse wave of amplitude  $y_m$  and wavelength  $\lambda_1$  is traveling on a stretched wire with tension  $\tau$ . The wave carry an average power  $P_{1\text{-avg}} = 0.48 \text{ W}$ . Then the wavelength of the wave is doubled ( $\lambda_2 = 2\lambda_1$ ) while keeping the tension  $\tau$  and amplitude  $y_m$  constant. What is the new average power  $P_{2\text{-avg}}$  carried by the wave with wavelength  $\lambda_2$ ?

- A) 0.12 W
- B) 0.44 W
- C) 0.27 W
- D) 0.36 W
- E) 0.39 W

Q4.

A 0.550 m long string fixed at both ends is vibrating in its fundamental mode. The maximum transverse acceleration of a point at the middle of the string is  $8.40 \times 10^3 \text{ m/s}^2$  and the maximum transverse velocity is 3.80 m/s. What is the wave speed of the transverse traveling waves on this string?

- A) 387 m/s
- B) 155 m/s
- C) 272 m/s
- D) 299 m/s
- E) 422 m/s

Q5.

One of the harmonic frequencies of a pipe closed at one end is 550 Hz. If the next-highest harmonic frequency of the pipe is 650 Hz, what is the length of the pipe?

- A) 1.72 m
- B) 2.21 m
- C) 2.88 m
- D) 3.35 m
- E) 4.12 m

---

Q6.

A train approaches a mountain at a speed of 75.0 km/h. The train driver sounds a whistle that emits a frequency of 420 Hz. What will be the frequency of the echo that the train driver hears reflected from the mountain?

- A) 474 Hz
- B) 211 Hz
- C) 298 Hz
- D) 322 Hz
- E) 374 Hz

---

Q7.

A point sound source radiates sound isotropically in all directions in air. At a distance of 5.00 m from the source the sound intensity level is 52.0 dB. If the sound frequency is 587 Hz, what is the pressure amplitude at this distance? (Density of air = 1.21 kg/m<sup>3</sup>)


- A)  $1.14 \times 10^{-2}$  Pa
- B)  $2.22 \times 10^{-2}$  Pa
- C)  $3.13 \times 10^{-2}$  Pa
- D)  $4.14 \times 10^{-2}$  Pa
- E)  $5.55 \times 10^{-2}$  Pa

---

Q8.

Two in phase speakers are emitting sound waves of same frequency in a room, as shown in **Figure 1**. For what lowest frequency, the sound from the speakers produce destructive interference at the micro phone located at a distance of 5.00 m in front of one of the speaker?

Fig#



- A) 206 Hz
- B) 322 Hz
- C) 366 Hz
- D) 316 Hz
- E) 101 Hz

---

Q9.

The density of gasoline at  $0.00\text{ }^{\circ}\text{C}$  is  $730\text{ kg/m}^3$ . What will be the density of gasoline if its temperature is raised to  $30.0\text{ }^{\circ}\text{C}$ . Assume the coefficient of volume expansion of gasoline  $\beta = 0.950 \times 10^{-3}\text{ }/^{\circ}\text{C}$ .

- A)  $709\text{ kg/m}^3$
- B)  $525\text{ kg/m}^3$
- C)  $555\text{ kg/m}^3$
- D)  $630\text{ kg/m}^3$
- E)  $677\text{ kg/m}^3$

---

Q10.

A metal bar is used to conduct heat. When the temperature at one end of the bar is  $100^{\circ}\text{C}$  and at the other is  $20\text{ }^{\circ}\text{C}$ , heat is transferred at a rate of  $95\text{ J/s}$ . If the temperature of the hotter end is reduced to  $80^{\circ}\text{C}$ , what will be the rate of heat transfer?

- A)  $71\text{ J/s}$
- B)  $52\text{ J/s}$
- C)  $86\text{ J/s}$
- D)  $91\text{ J/s}$
- E)  $89\text{ J/s}$

---

Q11.


One kg of ice is mixed with one kg of water at  $10.0\text{ }^{\circ}\text{C}$ . When thermal equilibrium is reached, the mixture contains total  $2.00\text{ kg}$  of ice at  $0.00\text{ }^{\circ}\text{C}$ . Determine the initial temperature of the ice. The specific heat of ice  $2092\text{ J/kg }^{\circ}\text{C}$ .

- A)  $-179\text{ }^{\circ}\text{C}$
- B)  $-101\text{ }^{\circ}\text{C}$
- C)  $-121\text{ }^{\circ}\text{C}$
- D)  $-155\text{ }^{\circ}\text{C}$
- E)  $-191\text{ }^{\circ}\text{C}$

Q12.

**Figure 2** shows a  $pV$ -diagram for an ideal gas process  $a$  to  $b$  in which its absolute temperature at  $b$  is one-fourth of its absolute temperature at  $a$ . How many joules of work was done in this process?

Fig#



- A)  $-56.8$  J
- B)  $-85.6$  J
- C)  $-35.5$  J
- D)  $-44.6$  J
- E)  $-20.5$  J

Q13.


Suppose the pressure of an ideal monatomic gas is tripled while its volume is halved. What happens to the internal energy of the gas?

- A) It increases.
- B) It decreases.
- C) It stays the same, as the described changes do not affect the internal energy.
- D) This depends on the molecular weight of the gas involved, thus this is indeterminate.
- E) None of the given answers.

Q14.

One-third of a mole of a monoatomic ideal gas is taken along the path  $abc$  shown in **Figure 3**. How much heat is transferred in the process  $abc$ ?

Fig#



- A)  $3 \times 10^3$  J

- B)  $1 \times 10^3$  J  
 C)  $2 \times 10^3$  J  
 D)  $5 \times 10^3$  J  
 E)  $4 \times 10^3$  J

Q15.

During a thermal expansion process at constant pressure, 970 J of heat are added to 1.75 mol of an ideal gas to heat it from 10.0 °C to 25.0 °C. The gas does + 223 J of work during the expansion. Calculate  $\gamma$  for the gas.

- A) 1.30  
 B) 1.70  
 C) 1.40  
 D) 1.50  
 E) 1.10

Q16.


How much work is required to compress 5.00 mol of air at 20.0 °C and 1.00 atm to one-fifth of the original volume in an adiabatic process? Assume air behaves as a diatomic ideal gas?

- A) 27.5 kJ  
 B) 21.1 kJ  
 C) 22.5 kJ  
 D) 23.3 kJ  
 E) 25.5 kJ

Q17.

**Figure 4** shows the thermodynamic cycles of two heat engines. Determine ratio of the thermal efficiencies  $\epsilon_1$  of engine 1 and  $\epsilon_2$  of engine 2 ( $\epsilon_1 / \epsilon_2$ ) if the same amount of heat is added per cycle to each engine.

Fig#



- A) 2.0  
 B) 0.5  
 C) 1.0  
 D) 1.5  
 E) 1.3

Q18.

One hundred and twenty gram of water is heated to 95.0 °C and then cooled to room temperature at 17.0 °C. The cooling process is essentially isothermal for the air in the room. Calculate the total change in entropy of the water-air system while the water cools down, assuming that all the heat lost by the water goes into the air.

- A) +15.5 J/K
- B) -15.5 J/K
- C) +24.1 J/K
- D) -23.2 J/K
- E) +19.4 J/K

---

Q19.

A heat engine does 10.0 J of work and exhausts 15.0 J of waste heat during each cycle. If the cold-reservoir temperature is 20.0 °C, what is the minimum possible temperature in °C of the hot reservoir?

- A) 215 °C
- B) 107 °C
- C) 127 °C
- D) 147 °C
- E) 169 °C

---

Q20.

Fifty grams of water at 15.0 °C is placed in the freezer compartment of a refrigerator with a coefficient of performance of 4.00. How much work is supplied to the refrigerator to convert 50.0 g water at 15.0 °C into ice at 0.00 °C.?

- A)  $4.95 \times 10^3$  J
  - B)  $1.33 \times 10^3$  J
  - C)  $2.45 \times 10^3$  J
  - D)  $2.88 \times 10^3$  J
  - E)  $3.58 \times 10^3$  J
-

