

## Final Exam (T-002)

- Q1 Q0 The linear density of a vibrating string is 1 g/m.  
002Q0 A transverse wave is propagating on the string and  
17 Q0 is given by the equation:  
Q0  $y(x,t) = 2.0 \cdot \sin(x - 40 \cdot t)$ ,  
Q0 where  $x$  and  $y$  are in meters and  $t$  is in seconds. What  
Q0 is the tension in the string?  
Q0  
A1 1.6 N.  
A2 1.9 N.  
A3 0.9 N.  
A4 2.1 N.  
A5 5.2 N.  
Q0
- Q2 Q0 A stationary source emits a sound wave of frequency  $f$ .  
002Q0 If a man travels toward this stationary source with a  
18 Q0 speed twice the speed of sound, he will observe the  
Q0 emitted sound to have a frequency of:  
Q0  
A1  $3 \cdot f$ .  
A2  $f/2$ .  
A3  $2 \cdot f$ .  
A4  $f$ .  
A5 indefinite frequency.  
Q0
- Q3 Q0 Which of the following statements are CORRECT:  
002Q0  
Q0 1. Waves carry energy and momentum.  
Q0 2. Mechanical waves need a medium to propagate.  
Q0 3. Sound waves are transverse waves.  
Q0 4. A Wave on a stretched string is a longitudinal wave.  
Q0 5. For a tube closed at one end, only odd harmonics are present.  
Q0  
A1 1, 2, and 5.  
A2 2 and 4.  
A3 1, 2 and 3.  
A4 1 and 4.  
A5 3 and 5.  
Q0
- Q4 Q0 3.00-kg of water at 100 degrees Celsius is converted to steam  
002Q0 at 100 degrees Celsius by boiling at one atmospheric pressure.  
20 Q0 For one kg of water, the volume changes from an initial value  
Q0 of  $1.0 \cdot 10^{(-3)} \text{ m}^3$  as a liquid to  $1.671 \text{ m}^3$  as steam.  
Q0 The work done by the water in this process is:  
Q0  
A1  $5.07 \cdot 10^{**5} \text{ J}$ .  
A2  $1.23 \cdot 10^{**5} \text{ J}$ .  
A3  $2.45 \cdot 10^{**5} \text{ J}$ .  
A4  $3.01 \cdot 10^{**5} \text{ J}$ .  
A5  $1.69 \cdot 10^{**5} \text{ J}$ .  
Q0
- Q5 Q0 The mass of a hydrogen molecule is  $3.3 \cdot 10^{**(-27)}$  kg. If  
002Q0  $1.0 \cdot 10^{**23}$  hydrogen molecules per second strike  $2.0 \text{ cm}^2$   
21 Q0 of wall at an angle of 55 degrees with the normal when  
Q0 moving with a speed of  $1.0 \cdot 10^{**3} \text{ m/s}$ , what pressure do

Q0 they exert on the wall?

Q0

A1  $1.9 \times 10^3$  Pa.

A2  $2.8 \times 10^3$  Pa.

A3  $5.7 \times 10^3$  Pa.

A4  $8.6 \times 10^3$  Pa.

A5  $0.9 \times 10^3$  Pa.

Q0

Q6 Q0 Which of the following statements are CORRECT:

002Q0

Q0 1. Two objects are in thermal equilibrium if they have the same temperature.

Q0 2. In an isothermal process, the work done by an ideal gas is equal to the heat energy

Q0 3. In an adiabatic process, no heat enters or leaves the system.

Q0 4. The thermal efficiency of an ideal engine can be = 1.0.

Q0 5. For any process the change in entropy of a closed system  $< 0$ .

Q0

A1 1, 2, and 3.

A2 4 and 5.

A3 1, 2 and 5.

A4 1 and 4.

A5 3 and 5.

Q0

Q7 Q0 Two positive charges,  $q_1$  and  $q_2$ , lie on the x-axis. The first charge,  $q_1 = 12.0 \times 10^{-6}$  C, is at the origin, and the second

002Q0 charge,  $q_2 = 3.0 \times 10^{-6}$  C, is at 3.0 m. Where must a

22 Q0 negative charge,  $q_3$ , be placed on the x-axis such that the resultant force on it is zero?

Q0

A1 2.0 m.

A2 - 1.5 m.

A3 3.0 m.

A4 1.0 m.

A5 - 1.0 m.

Q0

Q8 Q0 An electric dipole, of electric charge  $9.3 \times 10^{-12}$  C and distance  $1.0 \times 10^{-3}$  m, is in an electric field of strength

002Q0 1100 N/C. What is the difference in potential energy

23 Q0 corresponding to dipole orientations parallel and

Q0 anti-parallel to the field?

Q0

A1  $2.05 \times 10^{-11}$  J.

A2  $1.03 \times 10^{-11}$  J.

A3  $6.15 \times 10^{-15}$  J.

A4  $4.08 \times 10^{-13}$  J.

A5  $3.87 \times 10^{-11}$  J.

Q0

Q0

Q9 Q0 A ball of charge -50 e lies at the center of a hollow spherical metal shell that has a net charge of -100 e. What is the charge

002Q0 on the outer surface of the shell?

24 Q0

A1 -150 e.

A2 -100 e.

A3 - 50 e.

A4 100 e.

A5 150 e.

Q0

Q10Q0 In figure (8), a hollow sphere, of radius  $r$  that carries a  
25 Q0 negative charge  $-q$ , is put inside another hollow sphere, of  
002Q0 radius  $R$  that carries a positive charge  $Q$ . At a distance  $x$  from  
Q0 the common center, such that  $r < x < R$ , the potential is:

Q0

A1  $k \cdot [(Q/R) - (q/x)]$ .

A2  $k \cdot [(Q/R) - (q/r)]$ .

A3  $k \cdot [(Q/R) + (q/x)]$ .

A4  $k \cdot [(Q/R) + (q/r)]$ .

A5  $k \cdot [(Q/x) - (q/R)]$ .

Q0

Q0

Q11Q0 The electric potential in a certain region is given by:

25 Q0  $V(x,y,z) = -4x^2z - 5y + 3(z^2)$  Volts,

002Q0 where  $x$ ,  $y$  and  $z$  are in meters. What is the magnitude of  
Q0 the electric field at the point  $(+2, -1, -3)$ ?

Q0

A1 29 V/m.

A2 25 V/m.

A3 35 V/m.

A4 125 V/m.

A5 10 V/m.

Q0

Q12Q0 A parallel combination of two capacitors,  $C_1$  and  $C_2$

002Q0 where  $C_2 = 2C_1$ , is connected to a battery. If the charge

26 Q0 accumulated on  $C_1$  is  $2.0 \cdot 10^{(-6)}$  C and the total

Q0 energy stored in the combination is  $12.0 \cdot 10^{(-9)}$  Joule,

Q0 then the capacitance of  $C_2$  is:

Q0

A1  $1.0 \cdot 10^{(-3)}$  F.

A2  $2.5 \cdot 10^{(-6)}$  F.

A3  $1.5 \cdot 10^{(-6)}$  F.

A4  $1.5 \cdot 10^{(-3)}$  F.

A5  $3.0 \cdot 10^{(-6)}$  F.

Q0

Q13Q0 An electric device, which heats water by immersing a

27 Q0 resistance wire in the water, generates 153 J of heat per

992Q0 second when an electric potential difference of 12 V is

Q0 placed across its ends. What is the resistance of the

Q0 heater wire?

Q0

A1 0.94 Ohms

A2 0.81 Ohms

A3 0.58 Ohms

A4 0.48 Ohms

A5 2.10 Ohms

Q0

Q14Q0 The capacitor in figure (1) is initially charged to 50 V and

28 Q0 then the switch is closed. What charge flows out of the

002Q0 capacitor during the first minute after the switch was closed?

Q0

A1 4.8 mC.

A2 0.3 mC.

A3 3.6 mC.

A4 1.4 mC.

A5 1.7 mC.

Q15Q0

002Q0 In figure (2), if  $V_c - V_d = 6.0$  Volts, what is the emf of the  
28 Q0 battery?

Q0

A1 10.8 Volts.

A2 9.61 Volts.

A3 13.9 Volts.

A4 18.2 Volts.

A5 11.7 Volts.

Q0

Q16Q0 The sum of the currents entering a junction equals the sum  
28 Q0 of the currents leaving that junction is a consequence of:

991Q0

002A1 conservation of charge

A2 conservation of energy

A3 Coulomb's law

A4 Ampere's law

A5 Newton's second law

Q0

Q17Q0 Find the values of the currents in figure (3).

28 Q0

200Q0

A1  $I_1 = 2$  A,  $I_2 = 2$  A,  $I_3 = -4$  A.

A2  $I_1 = 2$  A,  $I_2 = 2$  A,  $I_3 = 4$  A.

A3  $I_1 = -2$  A,  $I_2 = -2$  A,  $I_3 = -4$  A.

A4  $I_1 = -2$  A,  $I_2 = 2$  A,  $I_3 = \text{zero}$ .

A5  $I_1 = 2$  A,  $I_2 = -2$  A,  $I_3 = \text{zero}$ .

Q0

Q0

Q18Q0 Which of the following statements are WRONG:

002Q0

Q0 1. In order to achieve the lowest resistance from several  
Q0 resistors, they should be connected in parallel.

Q0 2. In order to achieve the lowest capacitance from several  
Q0 capacitors, they should be connected in parallel.

Q0 3. The resistance of a conductor does not depend on temperature.

Q0 4. A dielectric increases the capacitance of a capacitor.

Q0 5. The electric flux through a closed surface is always zero.

Q0

A1 2, 3 and 5.

A2 2 and 4.

A3 1, 2 and 3.

A4 1 and 4.

A5 1 and 3.

Q0

Q19Q0 An electron is projected into a uniform magnetic field

29 Q0  $B = (0.8 \text{ k})$  T. Find the magnitude of the magnetic force,

002Q0 on the electron when the velocity is:

Q0  $v = (5.0 \cdot 10^{**5} \text{ i} + 3.0 \cdot 10^{**5} \text{ j})$  m/sec.

Q0 ( $\text{i}$ ,  $\text{j}$  and  $\text{k}$  are the unit vectors in the  $x$ ,  $y$  and  $z$  directions,  
Q0 respectively).

Q0

A1  $7.5 \cdot 10^{**(-14)}$  N.

A2  $5.2 \cdot 10^{**(-15)}$  N.

A3  $7.8 \cdot 10^{**(-18)}$  N.

A4  $1.2 \cdot 10^{**(-13)}$  N.

A5 ZERO.  
Q0

Q20Q0 In figure (4), a loop of wire carrying a current,  $I$ , of 2.0 A  
29 Q0 is in the shape of a right triangle with two equal sides, each  
002Q0 15 cm long. A 0.7 T uniform magnetic field is in the plane  
Q0 of the triangle and is perpendicular to the hypotenuse.  
Q0 The resultant magnetic force on the two equal sides is:  
Q0

A1 0.30 N, into the page.  
A2 0.30 N, out of the page.  
A3 0.41 N, into the page.  
A4 0.41 N, out of the page.  
A5 Zero.  
Q0

Q21Q0 A magnetic field CANNOT:  
29 Q0  
002A1 change the kinetic energy of a charge.  
A2 exert a force on a charge.  
A3 accelerate a charge.  
A5 exert a torque on a charged particle.  
A4 change the momentum of a charge.  
Q0

Q22Q0 Electrons are accelerated from rest through a potential  
29 Q0 difference of 500 V. They are then deflected by a magnetic  
993Q0 field of 0.2 T that is perpendicular to their velocity. The  
Q0 radius of the electrons trajectory is:  
Q0

A1 0.38 milli-m.  
A2 0.15 milli-m.  
A3 1.6 milli-m.  
A4 2.4 milli-m.  
A5 0.54 milli-m.  
Q0

Q23Q0 The current loop in figure (5) consists of one loop with two  
Q0 semicircles of different radii. If the current in the circuit  
29 Q0 is 19 A,  $a = 3.0$  cm and  $b = 5.0$  cm, then the magnetic dipole  
Q0 moment of the current loop is:  
Q0

A1  $0.10 \text{ A}\cdot\text{m}^2$ , into the page.  
A2  $0.10 \text{ A}\cdot\text{m}^2$ , out of the page.  
A3  $0.02 \text{ A}\cdot\text{m}^2$ , out of the page.  
A4  $0.02 \text{ A}\cdot\text{m}^2$ , into the page.  
A5  $1.15 \text{ A}\cdot\text{m}^2$ , into the page.  
Q0

Q24Q0 A conductor consists of a circular loop of radius  $R = 0.10$  m  
30 Q0 and two straight, long sections, as in Figure (6). The wire  
001Q0 lies in the plane of the paper (xy-plane) and carries a current  
Q0 of  $I = 5.3$  A. Determine the magnetic field, in Tesla, at the  
Q0 center of the loop. ( $\mathbf{k}$  is a unit vector in +z-direction)  
Q0

A1  $-4.4 \cdot 10^{(-5)} \text{ k}$ .  
A2  $5.8 \cdot 10^{(-5)} \text{ k}$ .  
A3  $-5.8 \cdot 10^{(-5)} \text{ k}$ .  
A4  $4.4 \cdot 10^{(-5)} \text{ k}$ .  
A5  $1.8 \cdot 10^{(-5)} \text{ k}$ .  
Q0

Q25Q0 A long solid cylindrical conductor of radius  $R = 4.0$  mm carries

30 Q0 a current  $I$  parallel to its axis. The current density in the  
993Q0 wire is  $2 \times 10^{**4}$  A/m\*\*2. Determine the magnitude of the magnetic  
Q0 field at a point that is 5.0 mm from the axis of the conductor.

- Q0
- A1 40 micro-T.
- A2 17 micro-T.
- A3 12 micro-T.
- A4 30 micro-T.
- A5 55 micro-T.

Q0  
Q26Q0 A solenoid is 3.0 m long and has a circumference of  
30 Q0  $9.4 \times 10^{**(-2)}$  m. It carries a current of 12.0 A. The magnetic  
Q0 field inside the solenoid is  $25.0 \times 10^{**(-3)}$  T. The length  
Q0 of the wire forming the solenoid is:

- Q0
- A1 467 m.
- A2 245 m.
- A3 233 m.
- A4 410 m.
- A5 900 m.

Q0  
Q0  
Q27Q0 Suppose that the identical currents  $I$  in figure (7) are all  
30 Q0 out of the page. The magnitude of the force per unit length  
002Q0 on the wire at the origin is:

- Q0 [take  $I = 10.0$  A, and  $a = 1.0 \times 10^{**(-4)}$  m.]
- Q0
- A1 0.28 N/m.
- A2 0.17 N/m.
- A3 0.18 N/m.
- A4 0.30 N/m.
- A5 0.55 N/m.

Q0  
Q28Q0 Faraday's law states that an induced emf is proportional to:

- 31 Q0
- 002Q0
- Q0
- A1 the rate of change of magnetic flux.
- A2 the rate of change of magnetic field.
- A3 the rate of change of electric flux.
- A4 the rate of change of electric field.
- A5 the rate of change of gravitational field.

Q0  
Q29Q0 A magnet is taken towards a metallic ring in such a way that  
31 Q0 a constant current of  $10^{**(-2)}$  A is induced in it. The total  
002Q0 resistance of the ring is 0.25 Ohm. In 10 seconds, the flux  
Q0 of the magnetic field through the ring changes by:

- Q0
- A1  $2.5 \times 10^{**(-2)}$  Wb.
- A2  $2.5 \times 10^{**(-3)}$  Wb.
- A3  $2.5 \times 10^{**(-6)}$  Wb.
- A4  $2.5 \times 10^{**(-1)}$  Wb.
- A5  $2.5 \times 10^{**(-9)}$  Wb.

Q0  
Q30Q0 Consider a circular loop of wire within which the magnetic flux,  
31 Q0  $\Phi$ , is given as a function of time,  $t$ , as

002Q0  $\Phi = a \cdot t^{**2} + b$ ,

Q0 where  $a$  and  $b$  are constants. If the induced emf is measured as  
Q0 48 V at  $t=3$  s, what is the value of  $a$ ?

Q0

A1 - 8.0 V/s.

A2 - 3.2 V/s.

A3 - 6.0 V/s.

A4 - 4.0 V/s.

A5 - 2.1 V/s.




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8