

Department of Physics, KFUPM

PHYSICS 102 – 053 – Final Exam – 17 August 17, 2006

Multiple Choice – (A) is the correct choice– ZERO VERSION

Q1.




Figure 1

Fig. 1 shows two graphs that represent a transverse wave on a string. Based on the information contained in these graphs, the speed of the wave is:

- A) 0.20 m/s.
- B) 0.30 m/s
- C) 0.40 m/s
- D) 0.10 m/s
- E) 0.80 m/s

Q2. As a sound wave travels from air into water, which of the following is TRUE?

- A) The frequency of the wave does not change
- B) The velocity of the wave decreases
- C) The wavelength of the wave decreases
- D) The wavelength of the wave does not change
- E) The frequency of the wave decreases

Q3. A transverse wave is traveling on a string. The displacement  $y$  of a particle on the string from its equilibrium position is given by  $y = 0.021\sin(2.0x - 25t)$ ,  $x$  and  $y$  are in meters, and  $t$  is in seconds. The linear density of the string is  $1.6 \times 10^{-2}$  kg/m. The tension in the string is

- A) 2.5 N
- B) 1.8 N
- C) 3.8 N
- D) 4.5 N
- E) 10 N

Q4. One cubic meter of water initially at 25 °C absorbs  $2.00 \times 10^8$  J of heat from the sun. Calculate the final temperature of the water. (Specific heat of water 4186 J/kg.K)

- A) 72.8 °C
- B) 92.5 °C
- C) -72.8 °C
- D) -92.8 °C
- E) 115 °C

Q5. A copper rod has one end in a heat reservoir of temperature 650 K and the other end at a heat reservoir of temperature 350 K. A total of 1200 J of heat flows from hot reservoir to cold reservoir through the rod. The total change in entropy of the two heat reservoirs is

- A) +1.6 J/K
- B) -1.6 J/K
- C) 3.5 J/K
- D) -3.5 J/K
- E) 0 J/K

Q6. A monatomic ideal gas is taken from A to B to C, as shown in Fig.2. The curved line between A and C is an isotherm. During the process the change in the internal energy





Figure 2

- A) zero
- B)  $8.0 \times 10^4$  J
- C)  $4.0 \times 10^4$  J
- D)  $-4.0 \times 10^4$  J
- E)  $-8.0 \times 10^4$  J

Q7. Two neutral metal spheres A and B on wood stands are touching (see Fig 3). A positively charged rod is held near sphere A but not touching it. While the rod is there, sphere B is moved so that the spheres no longer touch. Then the rod is removed. Afterward what is the charge state of each sphere?



- A) Sphere *A* negative, sphere *B* positive
- B) Sphere *A* positive, sphere *B* negative
- C) Sphere *A* neutral, sphere *B* negative
- D) Sphere *A* negative, sphere *B* neutral
- E) Sphere *A* neutral, Sphere *B* positive

Q8. A 0.100 g plastic sphere is charged by the addition of  $1.00 \times 10^{10}$  excess electrons. What electric field  $\vec{E}$  will cause the sphere to hang suspended in the air?

- A)  $6.13 \times 10^5$  N/C, vertically downward
- B)  $6.13 \times 10^5$  N/C, vertically upward
- C)  $-2.51 \times 10^5$  N/C, vertically downward
- D)  $2.51 \times 10^{+5}$  N/C, vertically upward
- E) 289 N/C, vertically downward

Q9. Two large and thin metal plates *A* and *B* are facing each other. The surface charge densities on the facing surfaces of the plates are  $+\sigma$  and  $-\sigma$  respectively and zero on the outer surfaces. Now plate *B* is removed very far from plate *A*. The charge density on plate *A* is:

- A)  $\frac{\sigma}{2}$
- B)  $\sigma$
- C)  $2\sigma$
- D)  $-\sigma$
- E) zero

Q10. A ball of radius 20 cm is uniformly charged to 80 nC. The magnitude of electric field strength at  $r = 10$  cm is

- A) 9000 N/C
- B) 18000 N/C
- C) 3000 N/C
- D) 36000 N/C
- E) 45000 N/C

Q11. The two segments of the wire in Fig. 4 have equal diameters but different resistivities  $\rho_1$  and  $\rho_2$ . Current  $I$  passes through this wire. If  $\rho_2/\rho_1 = \frac{1}{2}$ , what is the ratio of  $E_2/E_1$  of the electric field strengths in the two segments?




Figure 4

- A)  $\frac{1}{2}$
- B) 2
- C) 1
- D) 4
- E)  $\frac{1}{4}$

Q12. A proton's speed as it passes point A is  $5.0 \times 10^4$  m/s. It follows the trajectory shown in Fig. 5. What is the proton's speed at point B? (mass of the proton is  $1.67 \times 10^{-27}$  kg)




Figure 5

- A)  $1.2 \times 10^5$  m/s
- B)  $3.5 \times 10^5$  m/s
- C)  $4.0 \times 10^4$  m/s
- D)  $2.1 \times 10^6$  m/s
- E) zero

Q13. A battery with an emf of 60V is connected to the two capacitors shown in Fig. 6. The final charge on capacitor  $C_2$  is  $450 \mu\text{C}$ . What is the capacitance  $C_2$ ?




Figure 6

- A)  $20 \mu\text{F}$
- B)  $10 \mu\text{F}$
- C)  $30 \mu\text{F}$
- D)  $40 \mu\text{F}$
- E)  $5 \mu\text{F}$

Q14. In Fig. 7, what is the rate at which energy is supplied by the battery  $\mathcal{E}_1$  ?




Figure 7

- A)  $7.5 \text{ W}$
- B)  $2.1 \text{ W}$
- C)  $11.0 \text{ W}$
- D)  $20.1 \text{ W}$
- E)  $22.3 \text{ W}$

Q15. In Fig. 8, all the batteries are ideal with  $\mathcal{E}_1=6.0 \text{ V}$ ,  $\mathcal{E}_2=5.0 \text{ V}$ , and  $\mathcal{E}_3= 4.0 \text{ V}$ . What is the potential difference across resistor  $R_2$ ?




Figure 8

- A) 3 V
- B) 6 V
- C) 9 V
- D) 1.5 V
- E) 4.5 V

Q16. In Fig. 9,  $\mathcal{E} = 4.2 \text{ kV}$ ,  $C = 6.5 \text{ } \mu\text{F}$ ,  $R_1 = R_2 = R_3 = 0.92 \text{ M}\Omega$ . After switch  $S_1$  has been closed for a long time, what is the current in  $R_2$ ?




Figure 9

- A) 2.3 mA
- B) 4.6 mA
- C) 11 mA
- D) 8.2 mA
- E) zero

Q17. A battery has an emf of 12.00 volts. When a current  $I = 1.00 \text{ A}$  flows through the battery, the terminal voltage is 11.99 volts. What is the internal resistance of the battery?

- A)  $0.01 \text{ } \Omega$
- B)  $2.0 \text{ } \Omega$
- C)  $0.003 \text{ } \Omega$
- D)  $0.02 \text{ } \Omega$
- E)  $8.0 \text{ } \Omega$

Q18. A 22-V battery is connected across the terminals *a* and *b* in Fig. 10. If each resistor is  $40\ \Omega$ , what is the potential drop across the resistor labeled *R*?




Figure 10

- A) 8 V
- B) 11 V
- C) 14.7 V
- D) 12 V
- E) 16 V

Q19. The magnetic force on a point charge in a magnetic field is largest for a given speed when it:

- A) moves perpendicular to the magnetic field
- B) moves in the direction of the magnetic field
- C) moves in the direction opposite to the magnetic field
- D) has velocity components both parallel to and perpendicular to the field
- E) has velocity components both perpendicular and anti-parallel to the field.

Q20.




Figure 11

The parallel plates shown in Fig. 11 are 3.8 cm apart. A 0.064-T magnetic field is present in the space between the plates perpendicular to the plane of the paper. When an electron traveling horizontally with a speed of  $5.1 \times 10^5\ \text{m/s}$  enters the region, it passes through undeflected. The potential difference between the plates is:

- A) 1.24 kV

- B) 3.14 kV
- C) 10.1 kV
- D) 14.0 kV
- E) zero

Q21. A 62.8-m wire is made into a closely packed solenoid of diameter 1.00 cm. The length of the solenoid is 20.0 cm. What current through the wire will produce a magnetic field of 0.0126 T at its center?

- A) 1.00 A
- B) 8.21 A
- C) 2.31 A
- D) 4.21 A
- E) 3.11 A

Q22.




Figure 12

In Fig. 12, two circular arcs having radii  $a = 13.5$  cm and  $b = 10.7$  cm carry the same current  $i = 0.411$  A and share the same center of curvature P. The magnitude of the magnetic field at P is:

- A)  $1.08 \times 10^{-6}$  T
- B)  $8.41 \times 10^{-6}$  T
- C)  $2.11 \times 10^{-6}$  T
- D)  $9.89 \times 10^{-6}$  T
- E)  $1.11 \times 10^{-6}$  T

Q23. In Fig. 13, two long straight wires are perpendicular to the page. Each carries a current of 25.0 A directed out of the page. In unit vector notation, what is the net magnetic force per unit length on the wire at the origin?






Figure 13

- A)  $6.25 \times 10^{-4} (\hat{i} + \hat{j}) \text{ N/m.}$
- B)  $1.84 \times 10^{-4} (\hat{i} - \hat{j}) \text{ N/m.}$
- C)  $6.25 \times 10^{-4} (-\hat{i} - \hat{j}) \text{ N/m.}$
- D)  $1.84 \times 10^{-4} (-\hat{i} - \hat{j}) \text{ N/m.}$
- E)  $2.16 \times 10^{-4} (-\hat{i} + \hat{j}) \text{ N/m.}$

Q24. Fig. 14 shows the cross section of a long solid wire carrying a uniform current  $i$ . The radius of the wire is  $R$ . What is the value of the integral  $\oint \vec{B} \cdot d\vec{s}$  over the circular closed path of radius  $r$  shown in the Figure?




Figure 14

- A)  $\mu_0 i \left( \frac{r}{R} \right)^2$
- B)  $\mu_0 i \frac{r}{R^2}$
- C)  $\mu_0 i \frac{R}{r}$
- D)  $\mu_0 i \frac{R^2}{r}$
- E)  $\mu_0 i$

Q25.




Figure 15

In Fig. 15, the magnetic flux through the loop increases according to the relation  $\Phi_B = 2.0t^6 + 7$ , where  $\Phi_B$  is  $\text{T}\cdot\text{m}^2$  and  $t$  in seconds. The magnitude and direction of the current through the resistor  $R = 24 \Omega$  at  $t = 1 \text{ s}$  are:

- A) 0.50 A, counter clockwise
- B) 0.50 A, clockwise
- C) 1.5 A, counter clockwise
- D) 1.5 A, clockwise
- E) 2.8 A, counter clockwise

Q26. A wire of length 1.00 m is formed into a circular loop and placed perpendicular to a uniform magnetic field that is increasing at a constant rate of 20 mT/s. If the resistance of the wire is 100  $\Omega$ , at what rate is thermal energy generated in the loop?

- A)  $2.5 \times 10^{-8} \text{ W}$
- B)  $8.3 \times 10^{-8} \text{ W}$
- C)  $3.1 \times 10^{-8} \text{ W}$
- D)  $0.25 \times 10^{-8} \text{ W}$
- E)  $12 \times 10^{-8} \text{ W}$

Q27. The wing span (tip to tip) of a Boeing 747 airplane is 59 m. The plane is flying horizontally at a speed of 220 m/s. The vertical component of the earth's magnetic field is  $5.0 \times 10^{-5} \text{ T}$ . Find the induced emf between the wing tips.

- A) 0.65 V
- B) 0.032 V
- C) 2.5 V
- D) 0.12 V
- E) 1.8 V

Q28. In Fig. 16, a copper ring passes through a rectangular region where a constant magnetic field is directed into the page. In which position is the induced current through the ring is clockwise?




Figure 16

- A) 4
- B) 2
- C) 1
- D) 1 and 5
- E) 2,3 and 5

Q29. An ion of charge  $1.60 \times 10^{-19} \text{ C}$  has a mass of  $1.16 \times 10^{-26} \text{ kg}$ . It accelerates from rest through a potential of 500 V and enters a magnetic field of 0.400 T, moving perpendicular to the field. What is the radius of its circular path in the magnetic field?

- A) 2.13 cm
- B) 1.07 cm
- C) 4.19 cm
- D) 6.20 cm
- E) 12.5 cm

Q30. A certain coil of wire consists of 5 circular loops of radius 0.0400 m. It is placed in a region of uniform magnetic field parallel to the plane of the coil. The magnetic field is increasing at the rate of 0.200 T/s. The magnitude of the resulting induced emf is:

- A) zero
- B) 0.271 V
- C) 0.889 V
- D) 0.101 V
- E) 0.387 V