

Q1.

A standing wave having three nodes is set up in a string fixed at both ends. If the frequency of the wave is doubled, how many antinodes will there be?

- A) 4
- B) 3
- C) 2
- D) 5
- E) 6

Q2.

A stationary motion detector sends sound waves of frequency 0.120 MHz towards a truck approaching (the detector) at a speed of 50.0 m/s. What is the frequency of the waves reflected back to the detector? [Speed of sound = 343 m/s]

- A) 0.161 MHz
- B) 0.103 MHz
- C) 0.140 MHz
- D) 0.234 MHz
- E) 0.186 MHz

Q3.

A box with a total surface area of 1.20 m² and a wall thickness of 4.00 cm is made of an insulating material. A 10.0 W electric heater inside the box maintains the inside temperature at 15.0 °C above the outside temperature. Find the thermal conductivity of the insulating material.

- A) 0.022 W/m.K
- B) 2.20 W/m.K
- C) 0.034 W/m.K
- D) 0.016 W/m.K
- E) 1.23 W/m.K

Q4.

An ideal gas initially at 330 K is compressed at a constant pressure of 25.0 N/m² from a volume of 3.0 m³ to a volume of 1.00 m³. In the process, 75.0 J is lost by the gas as heat. What is the change in internal energy of the gas?

- A) -25.0 J
- B) -125 J
- C) +50.0 J
- D) +65.0 J
- E) -75.0 J

Q5.

A Carnot engine has an efficiency of ϵ when operating between $T_H = 400$ °C and $T_L = 200$ °C. What will be the efficiency of the same Carnot engine when operating between $T_H = 800$ °C and $T_L = 400$ °C.

- A) 1.25ϵ

- B) 2.0ϵ
- C) 0.50ϵ
- D) 1.5ϵ
- E) 0.77ϵ

Q6.

Two identical 0.20 kg masses are placed 1.0 m apart (center to center) on a frictionless surface. Each has $+10 \mu\text{C}$ of charge. What is the initial acceleration of one of the masses if it is released from rest and allowed to move?

- A) 4.5 m/s^2
- B) 2.3 m/s^2
- C) 6.3 m/s^2
- D) 1.4 m/s^2
- E) 5.2 m/s^2

Q7.


Two point charges $q_1 = +15 \mu\text{C}$ and $q_2 = -10 \mu\text{C}$ are placed in xy plane. If q_1 is placed at (20 cm, 0) and q_2 is placed at (0, 10 cm), find the resultant electric field at the origin.

- A) $(-3.38 \times 10^6 \hat{i} + 9.00 \times 10^6 \hat{j}) \text{ N/C}$
- B) $(3.38 \times 10^6 \hat{i} + 9.00 \times 10^6 \hat{j}) \text{ N/C}$
- C) $(-3.38 \times 10^6 \hat{i} - 9.00 \times 10^6 \hat{j}) \text{ N/C}$
- D) $(-9.00 \times 10^6 \hat{i} + 3.38 \times 10^6 \hat{j}) \text{ N/C}$
- E) $(9.00 \times 10^6 \hat{i} + 3.38 \times 10^6 \hat{j}) \text{ N/C}$

Q8.

FIGURE 1 shows portions of a large non-conducting sheet placed in parallel with long line of charge having a uniform charge per unit length λ . The surface charge density of the non-conducting sheet is $\sigma = +4.0 \mu\text{C}/\text{m}^2$. If the electric field intensity at point P, 2.0 cm on right of the line charge, is zero, find the linear charge density λ .

Fig#



- A) $-0.25 \mu\text{C}/\text{m}$
- B) $+0.25 \mu\text{C}/\text{m}$
- C) $+0.38 \mu\text{C}/\text{m}$
- D) $-0.38 \mu\text{C}/\text{m}$
- E) $-0.63 \mu\text{C}/\text{m}$

Q9.

A thick spherical conducting shell of outer radius 5.0 cm has a net charge $Q = +10 \mu\text{C}$. A point charge of $-3.0 \mu\text{C}$ is placed at its center. Find the surface charge density on the **outer surface** of the shell.

- A) $223 \mu\text{C}/\text{m}^2$
- B) $414 \mu\text{C}/\text{m}^2$
- C) $318 \mu\text{C}/\text{m}^2$
- D) $203 \mu\text{C}/\text{m}^2$
- E) $196 \mu\text{C}/\text{m}^2$

Q10.


Two conducting spherical shells of same outer radius $r = 5.0 \text{ cm}$ are placed at center to center distance of 20 cm. The charge of $+10 \mu\text{C}$ and $-20 \mu\text{C}$ is uniformly distributed over the outer surface of sphere 1 and 2, respectively, find the total potential at the center of sphere 1. (Assume potential is zero at infinity)

- A) +900 kV
- B) +270 kV
- C) -900 kV
- D) +450 kV
- E) +600 kV

Q11.

In **FIGURE 2** $C_1 = 2.0 \mu\text{F}$, $C_2 = 4.0 \mu\text{F}$, and $C_3 = 6.0 \mu\text{F}$. Find the equivalent capacitance between points a and b.

Fig#



- A) $8.7 \mu\text{F}$
- B) $18 \mu\text{F}$
- C) $9.8 \mu\text{F}$
- D) $12 \mu\text{F}$
- E) $2.0 \mu\text{F}$

Q12.


A current of 3.20 A exist in a copper wire whose diameter is 4.00 mm. The number of charge carrier per unit volume is $8.49 \times 10^{28} \text{ m}^{-3}$. Assuming the current density is uniform, calculate the electron drift speed.

- A) $1.87 \times 10^{-5} \text{ m/s}$
- B) $7.49 \times 10^{-5} \text{ m/s}$
- C) $4.25 \times 10^{-6} \text{ m/s}$
- D) $1.40 \times 10^{-5} \text{ m/s}$
- E) $3.23 \times 10^{-5} \text{ m/s}$

Q13.

Two real batteries with some internal resistances are shown in **FIGURE 3**. Find the potential difference between points A and B ($V_A - V_B$).

Fig#




- A) 13.0 V
- B) 23.5 V
- C) 11.8 V
- D) 24.5 V
- E) 18.0 V

Q14.

In a circuit shown in **FIGURE 4** if the potential of point **A** is zero find the potential of point **B**.

Fig#




- A) -2.0 V
- B) +2.0 V
- C) +6.0 V
- D) -6.0 V
- E) -4.0 V

Q15.

For the circuit given in **FIGURE 5**, if the current through one of the 6.00Ω is 0.500 A find the current through the 4.00Ω resistor.

Fig#




- A) 2.25 A
- B) 0.33 A
- C) 3.50 A
- D) 1.55 A
- E) 0.50 A

Q16.

If the current through the $150\ \Omega$ resistor is $1.00\ \text{A}$ as shown in **FIGURE 6**, find the current through the $51.0\ \Omega$ resistor.

Fig#




- A) 8.47 A
- B) 3.00 A
- C) 5.42 A
- D) 11.4 A
- E) 6.45 A

Q17.

Switch S in **FIGURE 7** is closed at time $t = 0$, to begin charging an initially uncharged capacitor $C = 15.0\ \mu\text{F}$ through a resistor $R = 20.0\ \Omega$. At what time is the potential difference across the capacitor double to that across the resistor?

Fig#




- A) $3.30 \times 10^{-4}\ \text{s}$
- B) $2.08 \times 10^{-4}\ \text{s}$
- C) $1.25 \times 10^{-4}\ \text{s}$
- D) $1.04 \times 10^{-4}\ \text{s}$
- E) $0.52 \times 10^{-4}\ \text{s}$

Q18.

A proton enters with the same speed in four regions of uniform magnetic fields of same magnitudes but in different directions, as shown in **FIGURE 8**. Rank regions according to magnitude of the force on the protons, **GREATEST FIRST**.

Fig#




- A) a and b tie, d, c
 B) b and c tie, a, d
 C) b and c tie, d, a
 D) a, then b and c tie, d
 E) a, d, b, c

Q19.

In **FIGURE 9** a charged particle electron or proton (you must decide which) moves in uniform magnetic field \vec{B} (out of the page), goes through half circle, and then exits that region. The charged particle spends 240 ns in the region, find the magnitude of magnetic field.

Fig#



- A) 0.137 T
 B) 0.298 T
 C) 1.49×10^{-4} T
 D) 3.72×10^{-4} T
 E) 0.579 T

Q20.

An 80.0 cm long wire, laying along the positive x axis (with one end at the origin), carries a current of 0.80 A in the negative x direction and placed in a magnetic field. $\vec{B} = 4.0\hat{i} + 12\hat{j}$, where x in meters and B in mT. Find, in unit vector notation, the magnetic force on the wire?


- A) $(-7.68 \times 10^{-3} \hat{k})N$
 B) $(+2.62 \times 10^{-3} \hat{k})N$
 C) $(-2.62 \times 10^{-3} \hat{k})N$
 D) $(+5.62 \times 10^{-3} \hat{j})N$

E) $(-4.72 \times 10^{-3} \hat{i})N$

Q21.

A rectangular loop with height $h = 6.50$ cm and width $w = 5.40$ cm is in a uniform magnetic field of magnitude $B = 0.250$ T, which points in negative y direction as shown in **FIGURE 10**. The loop makes an angle of $\theta = 30^\circ$ with the y axis and carries a current of 9.00 A in the direction indicated. What is the magnitude of the torque on the loop?

Fig#




- A) 6.84×10^{-3} N.m
- B) 9.56×10^{-3} N.m
- C) 7.62×10^{-3} N.m
- D) 3.95×10^{-3} N.m
- E) 4.32×10^{-3} N.m

Q22.

Two concentric circular loops of radii $r_1 = 20.0$ cm and $r_2 = 30.0$ cm are located in xy plane; each carries current of same magnitude $i = 7.00$ A but in opposite direction as shown in **FIGURE 11**. Find the net magnetic dipole moment of the loops.


Fig#



- A) $1.10 \text{ A}\cdot\text{m}^2$ into the page
- B) $2.86 \text{ A}\cdot\text{m}^2$ into the page
- C) $1.10 \text{ A}\cdot\text{m}^2$ out of the page
- D) $2.86 \text{ A}\cdot\text{m}^2$ out of the page
- E) $1.98 \text{ A}\cdot\text{m}^2$ into the page

Q23.

Wire 1 and wire 2 placed parallel to y axis at $x = 0$ and $x = 10 \text{ cm}$, respectively, and carries currents $i_1 = 4.0 \text{ A}$ and $i_2 = 8.0 \text{ A}$ in opposite directions as shown in **FIGURE 12**. A third wire carries current in positive y direction is to be placed parallel to wire 1 and wire 2 such that net force per unit length on wire 3 due to wire 1 and wire 2 is zero. Find the position x of wire 3.




- A) -10 cm
- B) $+20 \text{ cm}$
- C) $+15 \text{ cm}$
- D) -5.0 cm
- E) -15 cm

Q24.

In **FIGURE 13** wire 1 consist of a circular arc of radius R with central angle of $\theta = 120^\circ$ and two radial lengths and carries current $i_1 = 2.00 \text{ A}$ in the direction indicated. Wire 2 is long and straight and it carries a current i_2 and placed at a distance of $R/2$ from the center of circular arc. If the net magnetic field at the center of the arc O is zero find the current i_2 .

Fig#



- A) 1.05 A
- B) 2.75 A
- C) 0.84 A
- D) 1.34 A
- E) 2.05 A


Q25.

FIGURE 14 shows a cross section across the diameter of two long cylindrical conducting wires 1 and 2 of same radius R carrying same uniform current but in opposite directions. Four square paths (of side length R) of same dimensions are indicated for the line integral $\oint \vec{B} \cdot d\vec{s}$.

Rank the paths according to the magnitude of $\oint \vec{B} \cdot d\vec{s}$ taken in the directions shown,

GREATEST FIRST.

Fig#




- A) 4, then 1 and 3 tie, 2
- B) 1 and 3 tie, 2, 4
- C) 4, then 1, 2, and 3 tie
- D) 4, 3, 2, 1
- E) 3, 4, 1, 2

Q26.

A circular wire loop has radius of 0.12 m and carries current $i = 0.10\text{A}$ placed in the xy plane in a uniform magnetic field $\vec{B} = -1.5\hat{k}\text{T}$, as shown in **FIGURE 15**. Find the potential energy of the loop in the position shown.

Fig#



- A) $+6.79 \times 10^{-3}\text{ J}$
- B) $-6.79 \times 10^{-3}\text{ J}$
- C) $+5.65 \times 10^{-2}\text{ J}$
- D) 0
- E) $-5.65 \times 10^{-2}\text{ J}$

Q27.


A solenoid has a length $L = 1.55\text{ m}$ and an inner diameter $d = 4.15\text{ cm}$, and carries a current $i = 4.80\text{ A}$. The solenoid consists of six close-packed layers, each with 750 turns along length L . What is the magnitude of magnetic field at its center?

- A) 17.5 mT
- B) 23.8 mT
- C) 5.65 mT
- D) 13.5 mT
- E) 19.8 mT

Q28.

FIGURE 16 shows three situations in which a wire loop lies partially in a uniform magnetic field. The magnetic field is either increasing or decreasing, as indicated. In each situation, a battery is part of the loop. In which situation(s) is/are the induced emf and the battery emf in the same direction along the loop.

Fig#



- A) b only
B) b and c
C) a, b, and c
D) c only
E) a and c

Q29.

The magnetic flux through a loop increases according to the relation $\Phi_B = 6.0t^2 + 7.0t$, where Φ_B is in $\text{mT}\cdot\text{m}^2$ and t is in seconds. What is the magnitude of the emf induced in the loop when $t = 2.0$ s?

- A) 31 mV
B) 38 mV
C) 24 mV
D) 17 mV
E) 40 mV

Q30.

A 60.0 cm copper wire is formed into a square loop and placed perpendicular to a uniform magnetic field that is increasing at the constant rate 12.0 mT/s. If the resistance of the loop is 20.0 Ω , at what rate is thermal energy generated in the loop?

- A) 3.65×10^{-9} W
B) 2.70×10^{-9} W
C) 7.45×10^{-9} W
D) 1.35×10^{-9} W
E) 4.12×10^{-9} W