

Q1.

Standing waves pattern on a 6.00 m long string fixed at both ends is described by the wave function $y = 0.002 \sin(\pi x) \cos(100\pi t)$ where x and y are in meters and t is in seconds. How many loops are there in this standing wave pattern?

- A) 6
- B) 3
- C) 2
- D) 4
- E) 5

Q2.

A loudspeaker emits sound waves isotropically in all directions. What is the speaker's power output if the sound level is 90 dB at a distance of 20 m from the loud speaker?

- A) 5.0 W
- B) 3.5 W
- C) 1.5 W
- D) 4.0 W
- E) 2.5 W

Q3.

How many kg of ice at 0°C should be mixed with 1.8 kg of water at 80°C to bring the final temperature of the mixture to 10°C ?

- A) 1.4 kg
- B) 2.1 kg
- C) 2.4 kg
- D) 3.5 kg
- E) 1.1 kg

Q4.

2.00 L container of fixed volume holds 3.00 mol of an ideal gas. If 200 J of heat is added to the gas, what is the change in internal energy of the system?

- A) 200 J
- B) 150 J
- C) 100 J
- D) 170 J
- E) 110 J

Q5.

A monatomic ideal gas expands adiabatically from a volume of 2.0 liters to 6.0 liters. If the initial pressure is P_0 , what is the final pressure?

- A) $0.16 P_0$

- B) $9.0 P_0$
- C) $6.2 P_0$
- D) $3.0 P_0$
- E) $0.55 P_0$

Q6.

What is the change in entropy of 108 g of silver at a temperature of 961 °C when it is completely melted ($L_{F\text{-silver}} = 8.82 \times 10^4 \text{ J/kg}$, $T_{\text{Melting-silver}} = 961 \text{ °C}$).

- A) 7.72 J/K
- B) 5.53 J/K
- C) 3.21 J/K
- D) 1.33 J/K
- E) 6.11 J/K

Q7.


Coefficient of performance of an air conditioner is 2.80 and it operates on 800 W of power. Calculate the rate at which heat is discharged by the air conditioner to the outside air

- A) $3.04 \times 10^3 \text{ W}$
- B) $2.11 \times 10^3 \text{ W}$
- C) $1.35 \times 10^3 \text{ W}$
- D) $1.00 \times 10^3 \text{ W}$
- E) $4.35 \times 10^3 \text{ W}$

Q8.

Figure 1 shows two charged particles fixed on the x- axis. A third negatively charged particle can be placed at a certain point (1, 2, 3 or 4) on the x- axis so the net electrostatic force on it is zero. Which of the following answers can possibly be the correct position of the third particle?

Fig#



- A) 3
- B) 2
- C) 1
- D) 4
- E) None of the any given location

Q9.


An electron with a speed of 8.38×10^6 m/s enters a region of uniform electric field with velocity directed along the electric field. What is the magnitude of the electric field that will stop the electron momentarily at a distance of 0.100 m after entering this region?

- A) 2.00×10^3 N/C
- B) 1.14×10^3 N/C
- C) 1.32×10^3 N/C
- D) 2.42×10^3 N/C
- E) 1.22×10^3 N/C

Q10.

What is the magnitude of the electric flux through a horizontal surface of area 225 cm^2 placed in an electric field that makes 30.0° angle with the surface as shown in **Figure 2**?

Fig#




- A) $+2.00 \text{ N}\cdot\text{m}^2/\text{C}$
- B) $+3.55 \text{ N}\cdot\text{m}^2/\text{C}$
- C) $-1.12 \text{ N}\cdot\text{m}^2/\text{C}$
- D) $-2.00 \text{ N}\cdot\text{m}^2/\text{C}$
- E) $-3.55 \text{ N}\cdot\text{m}^2/\text{C}$

Q11.

Two particles of charges q_1 and q_2 are fixed in position, as shown in **Figure 3**. A third particle, of charge $+6.0 \mu\text{C}$, is brought from infinity to point P. Three particle system has the same electric potential energy as the initial two-particle system. What is the charge ratio q_1/q_2 ? (Assume potential energy is zero at infinity)

Fig#



- A) -1.4
- B) -1.1
- C) $+1.2$
- D) -1.8
- E) $+1.9$

Q12.


A solid conducting sphere of 10 cm radius has a net charge of 20 nC. If the potential at infinity is taken to be zero, what is the potential at the center of the sphere?

- A) 1.8×10^3 V
- B) 1.0×10^3 V
- C) 2.6×10^3 V
- D) 3.3×10^3 V
- E) Zero

Q13.

Six capacitors are connected in a circuit as shown in **Figure 4**. Find the energy stored in the equivalent capacitance of the circuit between points *a* and *b* if $C = 1.50 \mu\text{F}$ and the potential difference $V_{ab} = 100$ V.

Fig#



- A) 5.79×10^{-3} J
- B) 1.20×10^{-3} J
- C) 2.53×10^{-3} J
- D) 3.55×10^{-3} J
- E) 4.20×10^{-3} J

Q14.


Magnitude of the drift velocity of conduction electrons in a copper wire is 7.84×10^{-4} m/s and the number of conduction electrons per unit volume is $n = 8.46 \times 10^{28}$ /m³. What is the electric field in the wire? ($\rho_{\text{Copper}} = 1.72 \times 10^{-8}$ $\Omega \cdot \text{m}$)?

- A) 1.83×10^{-1} V/m
- B) 2.55×10^{-1} V/m
- C) 3.01×10^{-1} V/m
- D) 1.00×10^{-1} V/m
- E) 3.31×10^{-1} V/m

Q15.

In the circuit shown in **Figure 5**, a current of 0.25 A is flowing through the resistor R. What is the power dissipated in resistor R?

Fig#




- A) 0.38 W
 B) 0.55 W
 C) 0.11 W
 D) 0.73 W
 E) 0.92 W

Q16.

Figure 6 shows a circuit where the current in $2.0\ \Omega$ resistor is $3.0\ \text{A}$. Find the unknown emf \mathcal{E} .

Fig#




- A) 15 V
 B) 9.5 V
 C) 10 V
 D) 12 V
 E) 8.8 V

Q17.

For the circuit shown in **Figure 7**, find the potential difference $V_a - V_b$ across the $2.0\ \Omega$ resistor.

Fig#



- A) 8.0 V
B) 5.5 V
C) 9.1 V
D) 14 V
E) 10 V

Q18.

A capacitor is being charged through a $12\ \Omega$ resistor using a 10 V battery. What will be the current in the circuit when the capacitor has acquired $\frac{1}{4}$ of its maximum charge?

- A) 0.63 A
B) 0.42 A
C) 0.51 A
D) 0.29 A
E) 0.75 A

Q19.

A proton, enters a region of uniform magnetic field \vec{B} with a velocity $\vec{v} = 1.50\ \text{km/s}\hat{i}$. At that instant it experiences a magnetic force $\vec{F}_B = 2.25 \times 10^{-16}\ \text{N}\hat{j}$. What is the magnetic field \vec{B} ? Ignore the gravitational force.

- A) $-(0.938\ \text{T})\hat{k}$
B) $+(0.938\ \text{T})\hat{k}$
C) $-(0.532\ \text{T})\hat{k}$
D) $+(0.532\ \text{T})\hat{k}$
E) $-(0.232\ \text{T})\hat{k}$

Q20.

A proton moving in the positive x -direction with a speed $v = 1.35 \times 10^6\ \text{m/s}$ enters the region between the two plates as shown in **figure 8**. The potential of the top plate is 200 V, and the potential of the bottom plate is 0 V. What is magnetic field, \vec{B} , that is required between the plates so that the proton continues traveling in a straight line in the positive x -direction? Ignore the gravitational force.

Fig#



- A) $-(4.23 \times 10^{-3}\ \text{T})\hat{k}$
B) $+(4.23 \times 10^{-3}\ \text{T})\hat{k}$
C) $-(1.22 \times 10^{-3}\ \text{T})\hat{k}$
D) $+(1.22 \times 10^{-3}\ \text{T})\hat{k}$

E) $-(6.55 \times 10^{-3} \text{T}) \hat{k}$

Q21.


A charged particle undergoes uniform circular motion of radius $55.0 \mu\text{m}$ in a uniform magnetic field. The magnetic force on the particle has a magnitude of $2.80 \times 10^{-14} \text{ N}$. What is the kinetic energy of the particle?

- A) $7.70 \times 10^{-19} \text{ J}$
- B) $1.22 \times 10^{-19} \text{ J}$
- C) $2.56 \times 10^{-19} \text{ J}$
- D) $3.66 \times 10^{-19} \text{ J}$
- E) $5.34 \times 10^{-19} \text{ J}$

Q22.

A triangular loop of wire carrying a current of 0.125 A is placed in a x - y plane containing a uniform magnetic field $\vec{B} = 0.250 \text{ T} \hat{i}$, as shown in **Figure 9**. Determine the magnitude of the force on loop sides BC and CA, respectively due to the magnetic field.

Fig#



- A) $1.17 \times 10^{-3} \text{ N}$; $3.52 \times 10^{-3} \text{ N}$
- B) $1.47 \times 10^{-3} \text{ N}$; $2.22 \times 10^{-3} \text{ N}$
- C) $2.22 \times 10^{-3} \text{ N}$; $4.55 \times 10^{-3} \text{ N}$
- D) $3.52 \times 10^{-3} \text{ N}$; $4.22 \times 10^{-3} \text{ N}$
- E) $4.22 \times 10^{-3} \text{ N}$; $5.32 \times 10^{-3} \text{ N}$

Q23.


A circular loop of radius $r = 5.13 \text{ cm}$, has 47 turns. The loop is placed in a uniform magnetic field of magnitude 0.911 T . A current of 1.27 A flows through the loop. What is the maximum torque on the loop due to the magnetic field?

- A) 0.450 N.m
- B) 0.132 N.m
- C) 0.225 N.m
- D) 0.332 N.m
- E) 0.100 N.m

Q24.

Two long parallel wires, separated by a distance $D=10.0$ cm, each carry a current $I=5.00$ A, in opposite directions as shown in **Figure 10**. A circular loop, of radius $R = D/2$, has the same current I flowing in the counterclockwise direction. Determine the magnitude and the direction of the net magnetic field at the center of the loop P due to the current in the loop and in the parallel wires.

Fig#



- A) 1.03×10^{-4} T out of the page
 B) 1.03×10^{-4} T into the page
 C) 2.66×10^{-4} T out of the page
 D) 2.66×10^{-4} T into the page
 E) 3.45×10^{-4} T out of the page

Q25.


Two long parallel wires are separated by a distance of 3.0 mm. The current flowing in one of the wires is I and in the other wire is $2I$. If the magnitude of the force on a 1.0 m length of one of the wires is $7.0 \mu\text{N}$, what is the magnitude of current I ?

- A) 0.23 A
 B) 0.10 A
 C) 0.44 A
 D) 0.54 A
 E) 0.96 A

Q26.

A long solenoid with 6.00 cm diameter has 1000 turns per meter of thin wire which carries a current of 0.250 A. A long uniform straight wire carrying a current of 10.0 A is inserted along the axis of the solenoid, as shown in **Figure 11**. What is the magnitude of the magnetic field at a point 1.00 cm from the axis of the solenoid?

Fig#




- A) 3.72×10^{-4} T
 B) 1.00×10^{-4} T
 C) 1.52×10^{-4} T
 D) 2.11×10^{-4} T
 E) 2.44×10^{-4} T

Q27.

Figure 12 shows cross-sectional view of three wires that carry currents perpendicular to the plane the figure. The currents have magnitudes $I_1 = 3.0$ A, $I_2 = 4.0$ A and $I_3 = 4.0$ A in the directions shown. Four closed paths, labeled *a*, *b*, *c* and *d* are shown. Rank the magnitude of the line integral $\oint \vec{B} \cdot d\vec{l}$ for each path while going around the path in the counterclockwise direction, the **greatest first**.

Fig#




- A) *d, b, c, a*
 B) *a, b, c, d*
 C) *b, c, d, a*
 D) *c, d, a, b*
 E) *b, d, a, c*

Q28.

What is the net magnetic flux through the loop shown in **Figure 13**? Assume the area vector \vec{A} of the loop points into the page.

Fig#




- A) 4.0×10^{-2} Wb
 B) 1.0×10^{-2} Wb
 C) 1.5×10^{-2} Wb
 D) 2.1×10^{-2} Wb
 E) 2.7×10^{-2} Wb

Q29.

A circular wire loop has 4.8 cm radius and an electrical resistance of 0.16Ω . As shown in **Figure 14**, the loop is placed in a region where magnetic field \vec{B} is perpendicular to the loop. The magnetic field has an initial value of 8.0 T and is decreasing at a rate of 0.68 T/s. Determine the magnitude and direction of the induced current in the loop?

Fig#




- A) 3.1×10^{-2} A, counterclockwise
 B) 3.1×10^{-2} A, clockwise
 C) 1.9×10^{-2} A, counterclockwise
 D) 1.9×10^{-2} A, clockwise
 E) 1.0×10^{-2} A, clockwise

Q30.

A 10 cm long conducting rod moves at a constant speed $v = 0.50$ m/s on a zero-resistance horizontal wires towards 2.0Ω resistor in a uniform magnetic field $B = 0.50$ T, as shown in **Figure 15**. Find the magnitude of the force acting on the rod?

Fig#



- A) $6.3 \times 10^{-4}\text{ N}$
- B) $2.3 \times 10^{-4}\text{ N}$
- C) $1.3 \times 10^{-4}\text{ N}$
- D) $3.5 \times 10^{-4}\text{ N}$
- E) $4.4 \times 10^{-4}\text{ N}$