


Q1. In **FIGURE 1**, the three charges are:  $q_1 = +5.00 \mu\text{C}$ ,  $q_2 = +4.00 \mu\text{C}$ , and  $q_3 = -6.00 \mu\text{C}$ . What is the net electrostatic force on  $q_1$  due to  $q_2$  and  $q_3$ ?

Fig#




- A)  $-200 \hat{i} + 675 \hat{j}$  (N)
- B)  $-540 \hat{i} + 1350 \hat{j}$  (N)
- C)  $-120 \hat{i} + 380 \hat{j}$  (N)
- D)  $-222 \hat{i} + 750 \hat{j}$  (N)
- E)  $-133 \hat{i} + 300 \hat{j}$  (N)

Q2. Consider three distant spheres with charges  $Q_{1i} = 1\text{C}$ ,  $Q_{2i} = 2\text{C}$ , and  $Q_{3i} = 3\text{C}$ . We allow these three charges to touch each other for a short time and then we separate them. The new charges of these spheres become  $Q_{1f} = q$ ,  $Q_{2f} = 0.5q$ , and  $Q_{3f} = 1.5q$ . Find the value of  $q$ .

- A) 2 C
- B) 1 C
- C) 3 C
- D) 6 C
- E) 4 C

Q3. Particle 1 of charge  $Q_1 = 4Q$  and particle 2 of charge  $Q_2 = 9Q$  are fixed as shown in **FIGURE 2**. At what distance from  $Q_1$  along the y-axis will the net electric field due to the two particles be zero?


Fig#



- A)  $2L/5$
- B)  $4L/9$
- C)  $2L/3$
- D)  $3L/2$
- E)  $3L/5$

Q4. An electron is shot (ejected) at an initial speed of  $3.0 \times 10^4$  m/s at an angle of  $45^\circ$  relative to the positive x-axis, as shown in **FIGURE 3**. At time  $t = 0$ , the electron enters a region of uniform electric field  $\vec{E} = 2.0 \times 10^{-6} \hat{j}$  (N/C). Find the velocity of the electron along y-axis at  $t = 1.0$  s. Ignore gravity.


Fig#



- A)  $-3.2 \times 10^5 \hat{j}$  m/s
- B)  $+3.2 \times 10^5 \hat{j}$  m/s
- C)  $-2.5 \times 10^3 \hat{j}$  m/s
- D)  $+2.5 \times 10^3 \hat{j}$  m/s
- E)  $+3.2 \times 10^3 \hat{j}$  m/s

Q5. Consider three **different** electric dipoles placed in the same uniform electric field  $\vec{E}$ , as shown in **FIGURE 4**, where  $\theta = 60^\circ$ . Which of these dipoles has (or have) the **LOWEST** electric potential energy?


Fig#



- A) 2
- B) 1
- C) 3
- D) 1 and 3
- E) 2 and 3

Q6. A cube of 2.0 m edge is placed in a uniform field given by  $\vec{E} = 2.0\hat{i} + 1.0\hat{j}$  (N/C). The flux of the electric field through the face (**F**) perpendicular to the y- axis (see **FIGURE 5**) is


Fig#



- A)  $4.0 \text{ N}\cdot\text{m}^2/\text{C}$
- B)  $8.0 \text{ N}\cdot\text{m}^2/\text{C}$
- C)  $12 \text{ N}\cdot\text{m}^2/\text{C}$
- D)  $48 \text{ N}\cdot\text{m}^2/\text{C}$
- E) zero

Q7. In **FIGURE 6**, a charge  $q$  is placed at the common center of two hemispheres **A** and **B**. The flux of the electric field through hemisphere **B** is


Fig#



- A) equal to the flux through hemisphere **A**
- B) double the flux through hemisphere **A**
- C) four times the flux through hemisphere **A**
- D) zero
- E) half the flux through hemisphere **A**

Q8. Consider three infinite non-conducting sheets with uniform charge densities  $(-\sigma, +2\sigma, +3\sigma)$ , as shown in cross section in **FIGURE 7**. The electric field between plates **A** and **B** is given by

Fig#




- A)  $\frac{3\sigma}{\epsilon_0}$  to the left
- B)  $\frac{6\sigma}{\epsilon_0}$  to the left

- C)  $\frac{3\sigma}{\epsilon_0}$  to the right  
 D)  $\frac{6\sigma}{\epsilon_0}$  to the right  
 E)  $\frac{\sigma}{\epsilon_0}$  to the right

Q9. Two infinite wires are charged with uniform and opposite linear charge densities  $+\lambda$  and  $-\lambda$ , where  $\lambda = 1.00 \text{ nC/m}$ , as shown in **FIGURE 8**. The flux of the electric field through the Gaussian cylindrical surface (S) is


Fig#



- A)  $+2.26 \text{ N.m}^2/\text{C}$   
 B)  $-2.26 \text{ N.m}^2/\text{C}$   
 C)  $+113 \text{ N.m}^2/\text{C}$   
 D)  $-113 \text{ N.m}^2/\text{C}$   
 E) zero

Q10. A point charge  $q = -1.0 \times 10^{-10} \text{ C}$  is placed at the center of a spherical conducting shell that has a total charge  $Q = 5.0 \times 10^{-10} \text{ C}$ , as shown in **FIGURE 9**. The outer surface has radius  $R_2 = 10 \text{ cm}$ . The charge density on the external surface is equal to

Fig#



- A)  $+3.2 \text{ nC/m}^2$   
 B)  $-3.2 \text{ nC/m}^2$   
 C)  $+4.0 \text{ nC/m}^2$   
 D)  $+0.80 \text{ nC/m}^2$   
 E)  $-0.80 \text{ nC/m}^2$

Q11. A uniform electric field of magnitude 325 V/m is directed in the negative  $y$  direction. The coordinates of point A are  $(-0.200, -0.300)$  m, and those of point B are  $(0.400, 0.500)$  m. Calculate the electric potential difference  $V_B - V_A$ .


- A) +260 V
- B) -260 V
- C) -195 V
- D) +195 V
- E) +325 V

Q12. An electron is released from rest at the origin in a uniform electric field that points in the positive  $x$  direction and has a magnitude of 850 N/C. What is the change in the electric potential energy of the electron-field system when the electron moves a distance of 2.5 m?

- A)  $-3.4 \times 10^{-16}$  J
- B)  $+3.4 \times 10^{-16}$  J
- C)  $-1.4 \times 10^{-16}$  J
- D)  $+1.4 \times 10^{-16}$  J
- E)  $-5.4 \times 10^{-16}$  J

Q13. Two point charges ( $+q$  and  $-q$ ) are placed as shown in **FIGURE 10**. Consider the points 1, 2, 3, and 4 that are shown on the figure. At which point is the net electric potential **HIGHEST**? Take  $V = 0$  at infinity.


Fig#



- A) 4
- B) 3
- C) 2
- D) 1
- E) All points have the same potential

Q14. Four point charges are placed at the corners of a square, as shown in **FIGURE 11**. The magnitudes of the charges are equal. What is the electric potential energy of the system?

Fig#



- A) zero
- B)  $+5.4 \text{ kQ}^2/d$
- C)  $+2.6 \text{ kQ}^2/d$
- D)  $-2.6 \text{ kQ}^2/d$
- E)  $+0.71 \text{ kQ}^2/d$

Q15. In a certain region of space, the electric potential is given by:  $V = 2.0 \text{ } xyz^2$ , where  $V$  is in volts, and  $x$ ,  $y$ , and  $z$  are in meters. What is the magnitude of the electric field at the point with position vector  $(2.0\hat{i} - 2.0\hat{j} + 4.0\hat{k})$ ?


- A) 111 V/m
- B) 90.8 V/m
- C) 16.1 V/m
- D) 743 V/m
- E) 571 V/m

Q16. The electric potential immediately outside a charged conducting sphere is 200 V, and 10 cm farther from the surface of the sphere the potential is 150 V. What is the charge on the sphere?

- A) 6.7 nC
- B) 0.95 nC
- C) 1.3 nC
- D) 8.9 nC
- E) 5.4 nC

Q17. **FIGURE 12** shows a combination of four capacitors  $C_1 = C_3 = 8.0 \text{ } \mu\text{F}$  and  $C_2 = C_4 = 6.0 \text{ } \mu\text{F}$  connected to a 12-V battery. Calculate the charge on capacitor  $C_1$ .

Fig#



- A) 36  $\mu\text{C}$
- B) 18  $\mu\text{C}$
- C) 12  $\mu\text{C}$
- D) 24  $\mu\text{C}$
- E) 30  $\mu\text{C}$

Q18. A 20-V battery is connected to a series of  $N$  capacitors, each of capacitance  $4.0 \mu\text{F}$ . If the total energy stored in the capacitors is  $50 \mu\text{J}$ , what is  $N$ ?

- A) 16
- B) 12
- C) 4
- D) 10
- E) 8

Q19. A  $2\text{-}\mu\text{F}$  and a  $1\text{-}\mu\text{F}$  capacitor are connected in series and charged by a battery. They store charges  $P$  and  $Q$ , respectively. When disconnected and charged separately using the same battery, they have charges  $R$  and  $S$ , respectively. Then:

- A)  $R > S > P = Q$
- B)  $P > Q > R = S$
- C)  $R > P > S = Q$
- D)  $R > Q > S = P$
- E)  $S > R > P = Q$

Q20. A  $2.0\text{-nF}$  parallel plate capacitor is charged using a 12-V battery. The battery is removed and a dielectric of dielectric constant  $\kappa = 3.5$  is inserted, filling completely the space between the plates of the capacitor. What is the energy stored in the capacitor after inserting the dielectric?

- A)  $4.1 \times 10^{-8} \text{ J}$
- B)  $5.0 \times 10^{-5} \text{ J}$
- C)  $1.4 \times 10^{-7} \text{ J}$
- D)  $2.4 \times 10^{-8} \text{ J}$
- E)  $1.0 \times 10^{-6} \text{ J}$

**Physics 102**  
**Formula sheet for Second Major**

$F = \frac{kq_1q_2}{r^2}, \quad F = q_0 E$ $\Phi = \int_{\text{Surface}} \vec{E} \cdot d\vec{A}, \quad E = \frac{kq}{r^2}$ $\phi_c = \int \vec{E} \cdot d\vec{A} = \frac{q_{in}}{\epsilon_0}; \quad E = \frac{\sigma}{2\epsilon_0}; \quad E = \frac{\sigma}{\epsilon_0}$ $E = \frac{\sigma}{2\epsilon_0}; \quad E = \frac{\sigma}{\epsilon_0}$ $V = \frac{kQ}{r}, \quad W = \pm \Delta U$ $\Delta V = V_B - V_A = -\int_A^B \vec{E} \cdot d\vec{s} = \frac{\Delta U}{q_0}$ $E_x = -\frac{\partial V}{\partial x}, \quad E_y = -\frac{\partial V}{\partial y}, \quad E_z = -\frac{\partial V}{\partial z}$ $U = \frac{kq_1q_2}{r_{12}}$ $C = \frac{Q}{V}, \quad C_o = \frac{\epsilon_0 A}{d}, \quad C = 4\pi\epsilon_0 \frac{ab}{b-a},$ $U = -\vec{p} \cdot \vec{E} \quad \tau = \vec{p} \times \vec{E} \quad p = qd$	$v = v_o + at$ $x - x_o = v_o t + \frac{1}{2} at^2$ $v^2 = v_o^2 + 2a(x - x_o)$ <hr style="width: 50%; margin: 10px auto;"/> $\epsilon_0 = 8.85 \times 10^{-12} \text{ C}^2/\text{N.m}^2$ $k = 9.0 \times 10^9 \text{ N.m}^2/\text{C}^2$ $e = -1.6 \times 10^{-19} \text{ C}$ $m_e = 9.11 \times 10^{-31} \text{ kg}$ $m_p = 1.67 \times 10^{-27} \text{ kg}$ $g = 9.8 \text{ m/s}^2$ $\text{micro } (\mu) = 10^{-6}$ $\text{nano (n)} = 10^{-9}$ $\text{pico (p)} = 10^{-12}$
--	---