

Q1.

Two identical conducting spheres A and B carry equal charge  $Q$ , and are separated by a distance much larger than their diameters. Initially the electrostatic force between them is  $F$ . A third identical uncharged conducting sphere C is first touched to A, then to B, and then moved away. As a result of this, the electrostatic force between A and B becomes:

- A)  $3F/8$
- B)  $F/4$
- C)  $F/2$
- D)  $F/16$
- E)  $F$

Sec# Electric Charge - Coulomb's Law  
Grade# 44

Q2.

A positively charged sphere of mass 1.00 g falls from rest from a height of 5.00 m, in a uniform electric field of magnitude  $1.00 \times 10^4$  N/C and is directed vertically downward. The sphere hits the ground with a speed of 20.0 m/s. What is the charge on the sphere?


- A)  $+ 3.02 \mu\text{C}$
- B)  $- 1.00 \mu\text{C}$
- C)  $+ 5.23 \mu\text{C}$
- D)  $- 5.23 \mu\text{C}$
- E)  $+ 1.00 \mu\text{C}$

Sec# Electric fields - A point Charge in an Electric Field  
Grade# 47

Q3.

Figure 1 shows a dipole rotating under the effect of an electric field pointing along the negative x-axis. Which one of the following statements is TRUE

Fig#



- A) The potential energy of the dipole is decreasing.
- B) The torque on the dipole is directed into of the page.
- C) The dipole is rotating clockwise.
- D) The work done on the dipole by the field is negative.


E) The dipole will stop when it is pointing parallel to the positive x-axis.

Sec# Electric fields - A Dipole in an Electric Field  
Grade# 48

Q4.

Three point charges are located at the corners of a square as shown in Figure 2. Find the value of  $Q$  if the electric field at the corner A is zero. Take  $q = -7.00 \mu\text{C}$

Fig#




- A)  $19.8 \mu\text{C}$
- B)  $14.0 \mu\text{C}$
- C)  $9.90 \mu\text{C}$
- D)  $4.95 \mu\text{C}$
- E)  $2.54 \mu\text{C}$

Sec# Electric fields - The Electric Field Due to a Point Charge  
Grade# 54

Q5.

Figure 3 a, b and c, show the cross sections of three cylinders each carrying a uniform charge  $Q$ . Concentric with each cylinder is a cylindrical Gaussian surface, all three with the same radius. Rank the Gaussian surfaces according to the electric field at any point on the surface, GREATEST FIRST.

Fig#



- A) All tie
- B) a, b, c
- C) b, c, a
- D) c, b, a
- E) a, c, b

Sec# Gauss's law - Flux of an Electric Field  
Grade# 57

---

Q6.

A uniformly charged conducting sphere of 3.0 cm diameter has a surface charge density of  $10 \mu\text{C}/\text{m}^2$ . Find the total electric flux leaving the surface of the sphere.

- A)  $3.2 \times 10^3 \text{ N}\cdot\text{m}^2/\text{C}$
- B)  $1.3 \times 10^4 \text{ N}\cdot\text{m}^2/\text{C}$
- C)  $2.5 \times 10^3 \text{ N}\cdot\text{m}^2/\text{C}$
- D)  $1.4 \times 10^5 \text{ N}\cdot\text{m}^2/\text{C}$
- E)  $6.7 \times 10^2 \text{ N}\cdot\text{m}^2/\text{C}$

Sec# Gauss's law - Gauss's Law  
Grade# 55

---

Q7.

A  $6.0 \mu\text{C}$  charge is placed on a thin spherical conducting shell of radius  $R = 5.0 \text{ cm}$ . A particle with a charge of  $-10 \mu\text{C}$  is placed at the center of the shell. The magnitude and direction of the electric field at a point  $2R$  from the center of the shell are:

- A)  $3.6 \times 10^6 \text{ N/C}$ , toward the center
- B)  $3.6 \times 10^6 \text{ N/C}$ , away from the center
- C) 0
- D)  $5.4 \times 10^6 \text{ N/C}$ , toward the center
- E)  $5.4 \times 10^6 \text{ N/C}$ , away from the center

Sec# Gauss's law - A Charged Isolated Conductor  
Grade# 56

---

Q8.

A long, straight wire has fixed negative charge with a linear charge density of magnitude  $4.5 \text{ nC/m}$ . The wire is enclosed by a coaxial, thin walled nonconducting cylindrical shell of radius 20 cm. The shell is to have a positive charge on its outside surface (with a surface charge density  $\sigma$ ) that makes the net **electric** field at points 30 cm from the center of the shell equal to zero. Calculate  $\sigma$ .

- A)  $3.6 \times 10^{-9} \text{ C/m}^2$
- B)  $3.0 \times 10^{-10} \text{ C/m}^2$
- C)  $1.5 \times 10^{-10} \text{ C/m}^2$
- D)  $4.5 \times 10^{-7} \text{ C/m}^2$
- E)  $7.8 \times 10^{-5} \text{ C/m}^2$

Sec# Gauss's law - Applying Gauss's Law: Cylindrical Symmetry

Grade# 49

Q9.

Two large metal plates of area  $2.0 \text{ m}^2$  face each other,  $6.0 \text{ cm}$  apart, with equal charge magnitudes  $|q|$  but opposite signs. The magnitude of the electric field between the plates is  $1.2 \times 10^2 \text{ N/C}$ . Find  $|q|$ .

- A)  $2.1 \text{ nC}$
- B)  $1.1 \text{ nC}$
- C)  $0.50 \text{ nC}$
- D)  $13 \text{ nC}$
- E)  $0.40 \text{ nC}$

Sec# Gauss's law - Applying Gauss's Law: Planar Symmetry  
Grade# 60

Q10.

A glass sphere of diameter  $1.00 \text{ mm}$  has been charged to  $+100 \text{ nC}$ . A proton is fired from a large distance toward the sphere. What initial speed must the proton have to just reach the surface of the sphere? (Take  $V=0$  at a large distance from the sphere)


- A)  $1.86 \times 10^7 \text{ m/s}$
- B)  $9.10 \times 10^7 \text{ m/s}$
- C)  $5.34 \times 10^6 \text{ m/s}$
- D)  $4.50 \times 10^9 \text{ m/s}$
- E)  $2.67 \times 10^6 \text{ m/s}$

Sec# Electric Potential - Electric Potential Energy of a System of Point Charges  
Grade# 49

Q11.

Figure 4 shows a plot for the electric field  $E_x$  as a function of  $x$ . Find the magnitude of the potential difference between the points  $x = 2.00 \text{ m}$  and  $x = 6.00 \text{ m}$ .

Fig#



- A)  $14.5 \text{ V}$
- B)  $12.5 \text{ V}$
- C)  $10.0 \text{ V}$


- D) 16.5 V
- E) 11.0 V

Sec# Electric Potential - Potential Due to a Point Charge  
Grade# 46

Q12.

What are the magnitude and direction of the electric field at point P in Figure 5?

Fig#



- A)  $1.0 \times 10^4$  V/m to the left
- B)  $1.0 \times 10^4$  V/m to the right
- C)  $2.0 \times 10^4$  V/m to the left
- D)  $2.0 \times 10^4$  V/m to the right
- E)  $3.0 \times 10^3$  V/m upward

Sec# Electric Potential - calculating the Field from the Potential  
Grade# 46

Q13.

What is the charge on a conducting sphere of radius  $R = 0.20$  m if the potential at a distance  $r = 0.10$  m from the center of the sphere is 1500 V. (Take  $V = 0$  at infinity).


- A)  $3.3 \times 10^{-8}$  C
- B)  $1.7 \times 10^{-8}$  C
- C)  $1.5 \times 10^{-8}$  C
- D)  $2.5 \times 10^{-8}$  C
- E)  $4.5 \times 10^{-8}$  C

Sec# Electric Potential - Potential of a Charged Isolated Conductor  
Grade# 61

Q14.

In Figure 6, particles with charges  $q_1 = + 10 \mu\text{C}$  and  $q_2 = - 30 \mu\text{C}$  are fixed in place with a separation of  $d = 24$  cm. What is the value of  $Q$  that will make the potential equal zero at point P.

Fig#




- A)  $7.1 \mu\text{C}$
- B)  $5.1 \mu\text{C}$
- C)  $10 \mu\text{C}$
- D)  $3.5 \mu\text{C}$
- E)  $4.5 \mu\text{C}$

Sec# Electric Potential - Electric Potential Energy of a System of Point Charges  
Grade# 53

Q15.

Figure 7 shows three circuits, each consisting of a switch S and two capacitors, initially charged as indicated (top plate positive). After the switches have been closed, rank the charge on the right capacitor, GREATEST FIRST.

Fig#



- A) 1 and 2 tie, then 3
- B) 2, 1, 3
- C) All tie
- D) 3, 2, 1
- E) 3, 1, 2

Sec# Capacitance - Capacitors in Parallel and in Series  
Grade# 39

Q16.

Two capacitors are identical except that one is filled with air and the other is filled with oil. Both capacitors carry the same charge. If  $E_{\text{air}}$  refers to the electric field inside the capacitor filled with air, and  $E_{\text{oil}}$  refers to the electric field inside the capacitor filled with oil, then the ratio of the electric fields  $E_{\text{air}}/E_{\text{oil}}$  will be:


- A) greater than 1
- B) less than 1
- C) 0
- D) 1
- E) None of the other answers

Sec# Capacitance - Capacitor with a Dielectric  
Grade# 62

Q17.

Three identical capacitors are shown in Figure 8. A potential difference  $V = 10 \text{ kV}$  is established when the switch  $S$  is closed. Find the value of the capacitance  $C$  if the charge that passes through the meter  $M$  is  $0.20 \text{ C}$ .

Fig#




- A)  $6.7 \mu\text{F}$
- B)  $20 \mu\text{F}$
- C)  $1.6 \mu\text{F}$
- D)  $13 \mu\text{F}$
- E)  $2.5 \mu\text{F}$

Sec# Capacitance - Capacitors in Parallel and in Series  
Grade# 60

Q18.

Consider the circuit of identical capacitors shown in Figure 9. A potential difference of  $2.0 \times 10^2 \text{ V}$  is applied by the battery  $V$ . Calculate the energy stored in the system if the capacitance of each capacitor is  $50 \mu\text{F}$ .

Fig#



- A) 3.0 J
- B) 4.0 J
- C) 6.0 J
- D) 1.0 J
- E) 7.0 J

Sec# Capacitance - Energy stored in an Electric Field  
Grade# 55

Q19.

A cylindrical resistor of radius 2.5 mm and length 4.0 cm is made of a material that has a resistivity of  $3.5 \times 10^{-5} \Omega \cdot \text{m}$ . What is the potential difference when the energy dissipation rate in the resistor is 1.0 W?

- A) 0.27 V
- B) 1.8 V
- C) 2.2 V
- D) 0.17 V
- E) 1.1 V

Sec# Current and Resistance - Power in Electric Circuits  
Grade# 60

Q20.

A 1.0-m-long wire has a resistance equal to 0.30  $\Omega$ . A second wire made of identical material has a length of 2.0 m and a mass equal to the mass of the first wire. What is the resistance of the second wire?

- A) 1.2  $\Omega$
- B) 1.0  $\Omega$
- C) 3.4  $\Omega$
- D) 4.3  $\Omega$
- E) 5.6  $\Omega$



Sec# Current and Resistance - Ohm's Law  
Grade# 46

---

Test Expected Average = 52

---

---

$$F = k \frac{q_1 q_2}{r^2}$$

$$U = -\vec{p} \cdot \vec{E}$$

$$\vec{\tau} = \vec{p} \times \vec{E}$$

$$\Phi = \int_{\text{Surface}} \vec{E} \cdot d\vec{A}$$

$$\Phi_c = \iiint \vec{E} \cdot d\vec{A} = \frac{q_m}{\epsilon_0}$$

$$E = \frac{\sigma}{2\epsilon_0}$$

$$E = \frac{\sigma}{\epsilon_0}$$

$$E = k \frac{q}{r^2}$$

$$E = k \frac{q}{R^3} r$$

$$E = \frac{2k\lambda}{r}$$

$$\Delta V = V_B - V_A = -\int_A^B \vec{E} \cdot d\vec{S} = \frac{\Delta U}{q_0}$$

$$V = k \frac{q}{r}$$

$$E_x = -\frac{\partial V}{\partial x}, \quad E_y = -\frac{\partial V}{\partial y}, \quad E_z = -\frac{\partial V}{\partial z}$$

$$U = k \frac{q_1 q_2}{r_{12}}$$

$$C = \frac{q}{V}$$

$$C = \kappa C_{air}$$

$$U = \frac{1}{2} CV^2$$

$$U = \frac{1}{2} \epsilon_0 E^2$$

$$I = \frac{dQ}{dt}$$

$$I = JA$$

$$R = \frac{V}{I} = \rho \frac{L}{A}$$

$$J = \sigma E$$

$$\rho = \rho_0 [1 + \alpha(T - T_0)]$$

$$P = IV$$

---


$$v = v_0 + at$$

$$x - x_0 = v_0 t + \frac{1}{2} a t^2$$

$$v^2 = v_0^2 + 2 a (x - x_0)$$

### Constants:

$$k = 9.00 \times 10^9 \text{ N}\cdot\text{m}^2/\text{C}^2$$

$$\epsilon_0 = 8.85 \times 10^{-12} \text{ C}^2/\text{N}\cdot\text{m}^2$$

$$e = 1.60 \times 10^{-19} \text{ C}$$

$$m_e = 9.11 \times 10^{-31} \text{ kg}$$

$$m_p = 1.67 \times 10^{-27} \text{ kg}$$

$$g = 9.8 \text{ m/s}^2$$

---


$$\mu = \text{micro} = 10^{-6}$$

$$\text{n} = \text{nano} = 10^{-9}$$

$$\text{p} = \text{pico} = 10^{-12}$$