

Q1.

Two charges $q_1 = + 6.00 \mu\text{C}$ and $q_2 = -12.0 \mu\text{C}$ are placed at $(-2.00 \text{ cm}, 0)$ and $(4.00 \text{ cm}, 0)$, respectively. If a third unknown charge q_3 is to be located such that the net force on it from charges q_1 and q_2 is zero, what must be the coordinates of q_3 ?

- A) $(-16.5 \text{ cm}, 0)$
- B) $(-14.5 \text{ cm}, 0)$
- C) $(2.49 \text{ cm}, 0)$
- D) $(0, 0)$
- E) $(-6.50 \text{ cm}, 0)$

Q2.

Two small metallic spheres A and B carry $+ 1.00 \mu\text{C}$ and $-1.00 \mu\text{C}$ of charge, respectively, held fixed at a certain distance without touching each other. How many electrons must be transferred from one sphere to the other to reduce the force of attraction between them by a factor of four?

- A) 3.13×10^{12} from B to A
- B) 3.13×10^{12} from A to B
- C) 7.23×10^{12} from B to A
- D) 7.23×10^{12} from A to B
- E) 1.71×10^{11} from B to A

Q3.

The electric field midway between two charges of $+3q$ and $-2q$ is 98.0 N/C and the distance between the charges is 20.0 cm . What is the value of the charge q ?


- A) $2.18 \times 10^{-11} \text{ C}$
- B) $4.67 \times 10^{-6} \text{ C}$
- C) $1.09 \times 10^{-10} \text{ C}$
- D) $1.05 \times 10^{-5} \text{ C}$
- E) $5.73 \times 10^{-11} \text{ C}$

Q4.

FIGURE 1 shows four situations in which four charged particles are evenly spaced to the left and right of a central point P. The charge values are indicated. Rank the situations according to the magnitude of the net electric field at the central point, **GREATEST FIRST**.

Fig#

- A) 2, 4, 3, 1
- B) 4, 3, then 1 and 2 tie
- C) 3 and 4 tie, then 1 and 2 tie
- D) 4, 3, 1, 2



E) 1,4, 3, 2

Q5.

At some instant the velocity components of an electron moving between two parallel plates are $v_x = 1.5 \times 10^5 \text{ m/s}$ and $v_y = 3.0 \times 10^5 \text{ m/s}$. The uniform electric field between the plates is $\vec{E} = (1.2 \hat{j}) \text{ N/C}$. In unit vector notation find the velocity of the electron after $2.0 \mu\text{s}$. (ignore the effect of gravity)

- A) $(1.5 \times 10^5 \hat{i} - 1.2 \times 10^5 \hat{j}) \text{ m/s}$
- B) $(1.5 \times 10^5 \hat{i} + 7.2 \times 10^5 \hat{j}) \text{ m/s}$
- C) $(5.7 \times 10^5 \hat{i} - 1.2 \times 10^5 \hat{j}) \text{ m/s}$
- D) $(1.2 \times 10^5 \hat{i} - 1.2 \times 10^5 \hat{j}) \text{ m/s}$
- E) $(1.5 \times 10^5 \hat{i} + 3.0 \times 10^5 \hat{j}) \text{ m/s}$

Q6.

An electric dipole is placed in a uniform electric field $\vec{E} = (4000 \hat{i}) \text{ N/C}$. What is the change in dipole's potential energy if the initial and the final electric dipole moments \vec{p}_i and \vec{p}_f respectively, are given by

$$\vec{p}_i = (3.72 \times 10^{-30} \hat{i} + 4.96 \times 10^{-30} \hat{j}) \text{ C.m}$$


$$\vec{p}_f = (6.20 \times 10^{-30} \hat{i}) \text{ C.m}$$

- A) $-9.92 \times 10^{-27} \text{ J}$
- B) $+1.45 \times 10^{-27} \text{ J}$
- C) $+3.97 \times 10^{-26} \text{ J}$
- D) $+9.92 \times 10^{-27} \text{ J}$
- E) $-3.97 \times 10^{-26} \text{ J}$

Q7.

Consider a closed triangular box resting within a horizontal electric field of magnitude $E = 7.80 \times 10^4 \text{ N/C}$, as shown in **FIGURE 2**. Calculate the electric flux through the slanted surface.

Fig#




- A) $2.34 \times 10^3 \text{ N.m}^2/\text{C}$
- B) $1.17 \times 10^3 \text{ N.m}^2/\text{C}$

- C) $4.34 \times 10^3 \text{ N.m}^2/\text{C}$
- D) $7.34 \times 10^3 \text{ N.m}^2/\text{C}$
- E) $1.65 \times 10^3 \text{ N.m}^2/\text{C}$

Q8.

A long, straight wire has a linear charge density $\lambda = -7.8 \text{ nC/m}$. The wire is enclosed by a coaxial, thin-walled non-conducting cylindrical shell of radius 2.0 cm, as shown **FIGURE 3**. Find the surface charge density σ , on the outer surface of the shell that makes the net electric field zero outside the shell.

Fig#




- A) $6.21 \times 10^{-8} \text{ C/m}^2$
- B) $3.82 \times 10^{-8} \text{ C/m}^2$
- C) $1.24 \times 10^{-8} \text{ C/m}^2$
- D) $9.37 \times 10^{-8} \text{ C/m}^2$
- E) $5.31 \times 10^{-8} \text{ C/m}^2$

Q9.

FIGURE 4 shows, in cross section, three infinity large non-conducting sheets with uniform charge densities $\sigma_1 = +3.0 \mu\text{C/m}^2$, $\sigma_2 = +5.0 \mu\text{C/m}^2$, and $\sigma_3 = -7.0 \mu\text{C/m}^2$, and $L = 2.5 \text{ cm}$. In unit vector notation, what is the net electric field at point P?

Fig#




- A) $(+5.6 \times 10^4 \hat{j}) \text{ N/C}$
- B) $(-6.8 \times 10^5 \hat{j}) \text{ N/C}$
- C) $(-5.6 \times 10^4 \hat{j}) \text{ N/C}$
- D) $(+6.8 \times 10^5 \hat{j}) \text{ N/C}$
- E) $(-1.1 \times 10^5 \hat{j}) \text{ N/C}$

Q10.

A solid conducting sphere of radius a carries a net charge of $+2Q$. A conducting spherical shell of inner radius b and outer radius c is concentric with the solid sphere and carries a net charge $-Q$ as shown in **FIGURE 5**. Arrange the electric field in four regions labeled 1, 2, 3, and 4, **GREATEST FIRST**.

Fig#



- A) 2, 4 then 1 and 3 tie
- B) 1, 2, 3, 4
- C) 2, 3, 4, 1
- D) 1, 3, 2, 4
- E) 4 and 2 tie, then 3, 1

Q11.


Work done by the *electric field* in moving a charge $q = -4.0$ mC from point A to point B is $+20$ mJ. What is the potential difference between points A and B, i.e. $V_B - V_A = ?$

- A) $+5.0$ V
- B) -5.0 V
- C) -20 V
- D) $+20$ V
- E) -4.0 V

Q12.

FIGURE 6 shows four arrangements of charged particles (where $q > 0$), all located at the same distance from the origin. Rank the situations according to the net electric potential at the origin, **MOST POSITIVE FIRST** (take the potential to be zero at infinity).

Fig#



- A) b then a, c, and d all tie

- B) d, a, b, c
- C) a and b tie then c and d tie
- D) d, c, a, b
- E) c, b, a, d

Q13.


The electric potential in xy plane is given by $V = (2.0 x^2 - 3.0 y^2)$, where V is in volts and x and y are in meters. In unit vector notation, what is the electric field at point $(5.0m, 7.0m)$?

- A) $(-20\hat{i} + 42\hat{j})V/m$
- B) $(+20\hat{i} - 42\hat{j})V/m$
- C) $(-50\hat{i} + 95\hat{j})V/m$
- D) $(-20\hat{i} + 95\hat{j})V/m$
- E) $(-30\hat{i} + 42\hat{j})V/m$

Q14.

Two charges each of $+2.0 \mu\text{C}$ are fixed along the x axis at $x = -0.40 \text{ m}$ and $x = +0.40 \text{ m}$, as shown in **FIGURE 7**. A third charge $q = +4.0 \mu\text{C}$ is released from rest from $y = 0.3 \text{ m}$. Find the maximum kinetic energy attained by the third charge after the release.

Fig#



- A) 0.29 J
- B) 0.54 J
- C) 0.24 J
- D) 0.73 J
- E) 0.13 J

Q15.


An air filled parallel plate capacitor is charged to a potential difference of 10 V and then disconnected from the battery. Now if you fill the space between the plates with dielectric material of $k = 2$, which of the following statements is correct

- A) The voltage across the capacitor reduces to 5.0 V.
- B) The voltage across the capacitor remains at 10 V.
- C) The capacitance remains the same.
- D) The energy stored by the capacitor remains the same.
- E) The charge stored reduces to half of its initial value.

Q16.

If $C = 50.0 \mu\text{F}$ in **FIGURE 8**, what is the equivalent capacitance between points A and D?

Fig#




- A) $42.3 \mu\text{F}$
- B) $325 \mu\text{F}$
- C) $59.1 \mu\text{F}$
- D) $28.3 \mu\text{F}$
- E) $13.7 \mu\text{F}$

Q17.

In **FIGURE 9**, the capacitors $C_1 = 2.0 \mu\text{F}$ and $C_2 = 4.0 \mu\text{F}$. The capacitor C_1 is charged to a potential difference of 10 V and C_2 is initially uncharged. After closing the switch S , find the total energy stored by the two capacitors.

Fig#




- A) $3.3 \times 10^{-5} \text{ J}$
- B) $1.0 \times 10^{-4} \text{ J}$
- C) $3.0 \times 10^{-4} \text{ J}$
- D) $1.5 \times 10^{-5} \text{ J}$
- E) $5.0 \times 10^{-5} \text{ J}$

Q18.

An air filled parallel plate capacitor has capacitance of $C = 20.0 \mu\text{F}$. If lower half of the capacitor is filled with dielectric material $\kappa = 10$, as shown in **FIGURE 10**, find the capacitance of the new capacitor.

Fig#



- A) 110 μF
- B) 9.09 μF
- C) 440 μF
- D) 73.5 μF
- E) 173 μF

Q19.

When 115 V is applied across a 10 m long wire with a 0.30 mm radius, the magnitude of the current density is $1.4 \times 10^8 \text{ A/m}^2$. Find the resistivity of the wire.

- A) $8.21 \times 10^{-8} \Omega\cdot\text{m}$
- B) $9.33 \times 10^{-7} \Omega\cdot\text{m}$
- C) $5.72 \times 10^{-8} \Omega\cdot\text{m}$
- D) $2.38 \times 10^{-8} \Omega\cdot\text{m}$
- E) $4.16 \times 10^{-8} \Omega\cdot\text{m}$

Q20.

A heater element with potential difference of 400 V across it transfers electrical energy to thermal energy at the rate of 3000 W. At what rate the heater will transfer the energy if it is connected across a potential difference 300 V.

- A) 1.69 kW
- B) 2.25 kW
- C) 3.57 kW
- D) 1.13 kW
- E) 2.92 kW